

A LIST OF THE TYPE-SPECIMENS OF
LIBELLULIDAE AND CORDULIIDAE
(ODONATA) IN THE BRITISH MUSEUM
(NATURAL HISTORY)



BY

D. E. KIMMINS

British Museum (Natural History)

Pp. 277-305

BULLETIN OF
THE BRITISH MUSEUM (NATURAL HISTORY)
ENTOMOLOGY

Vol. 22 No. 6

LONDON: 1968

THE BULLETIN OF THE BRITISH MUSEUM (NATURAL HISTORY), instituted in 1949, is issued in five series corresponding to the Departments of the Museum, and an Historical series.

Parts will appear at irregular intervals as they become ready. Volumes will contain about three or four hundred pages, and will not necessarily be completed within one calendar year.

In 1965 a separate supplementary series of longer papers was instituted, numbered serially for each Department.

This paper is Vol. 22, No. 6 of the Entomological series. The abbreviated titles of periodicals cited follow those of the World List of Scientific Periodicals.

World List abbreviation :
Bull. Br. Mus. nat. Hist. (Ent.).

© Trustees of the British Museum (Natural History) 1968

TRUSTEES OF
THE BRITISH MUSEUM (NATURAL HISTORY)

Issued 20 September, 1968

Price Twelve Shillings.

A LIST OF THE TYPE-SPECIMENS OF LIBELLULIDAE AND CORDULIIDAE (ODONATA) IN THE BRITISH MUSEUM (NATURAL HISTORY)

By D. E. KIMMINS

SYNOPSIS

A list of the Libellulid and Corduliid type-specimens in the British Museum (Nat. Hist.) has been prepared. Three hundred and forty-three taxa are dealt with and lectotypes are designated for ninety-seven of these taxa.

INTRODUCTION

IN 1966, following the bequest to the British Museum (Nat. Hist.) of the Fraser collection of Odonata, the author published a list of Fraser's Odonata types in the BM(NH) collections. This produced valuable information and the present work offers a similar list of the type-specimens of Libellulidae and Corduliidae in our collection. The publication of such lists is advocated by the International Commission on Zoological Nomenclature (Rec. 72D(4)) and the preparation of this list has proved both useful and interesting. Certain discrepancies in the labelling and marking of some of the earlier types have come to light and the opportunity has been taken to designate lectotypes in ninety-three cases in which the descriptions were based upon a series of specimens and a type or holotype was not specified by the original author in print. The word LECTOTYPE, in capital letters, indicates a designation of lectotype in this paper.

This list includes not only Fraser's types, dealt with by Kimmins (1966), but also the small number of Fabrician Libellulid types from the Sir Joseph Banks collection, which were studied by Campion (1917). In both these cases, the original type data is not repeated, but reference is made to the appropriate paper in which it may be found.

The general lay-out follows that of Kimmins (1966), each entry beginning with the specific name (in alphabetical order), author, the genus in which it was described, followed by the date and page and figure references. Full references are given at the end of the paper. Next follows the status of the type specimen and the label data, that of each label being separated from the next by an oblique stroke.

Any comments are given in a second paragraph. Square brackets enclose additional information, such as the completion of abbreviated locality names; [WFK] indicates that a label is in Kirby's writing. Type [McL. label] indicates the small, rectangular type-label, usually on mauve or reddish paper, used by McLachlan. BM Register numbers are not given.

In the case of species described by Calvert (1909), it appears to have been his practice to label as TYPE the example figured. These are to be considered as Holotypes. (Calvert, 1901-08: xxviii).

This list of types was completed in February 1968.

adolescens Kirby (*Beblecia*), 1900 : 71-72, fig. 2. LECTOTYPE ♂ (abdominal segments 5-10 missing). Free Town, Sierra Leone, 13.ix.99, *E. E. Austen*/*Beblecia adolescens* Kb. type [WFK].

Of the three examples listed by Kirby, only one (the lectotype) bears his determination label, as recorded above, but the other two examples, 1 ♂, 1 ♀, are recognizable by the locality data and are considered to be paralectotypes. No type was designated in print, but the example bearing his determination label has been selected as Lectotype. This specimen has for many years carried a red type label, now replaced by a lectotype label. This taxon is currently considered a synonym of *Crocothemis divisa* Karsch.

aenea Kirby (*Zygonidia*), 1905 : 275-277. Holotype ♂. Tonkin/*Zygonidia aenea* Kirby, ♂ Holotype, D. E. Kimmins det. 1961.

Currently placed as a synonym of *Zygonyx iris* Selys.

aequalis Kimmins (ssp. *Rhyothemis phyllis*), 1936 : 78-80, text-fig. 7 and pl. 3. Holotype ♂. New Hebrides, Malekula, Ounua, 21.iii.1929, Miss L. E. Cheesman/*Rhyothemis phyllis* aequalis Kimmins ♂, det. D. E. Kimmins.

aethiopica Kimmins (ssp. *Atoconeura biordinata*), 1958 : 357-358, figs. 7a-c. Holotype ♂. Ethiopia, Segheria, 25.iii.1948, K. M. Guichard/*Atoconeura biordinata aethiopica* Kim., ♂ Type, D. E. Kimmins det. 1958.

affinis Kirby (*Neurothemis*), 1889 : 323, pl. 54, fig. 2. Holotype ♂. Barang/N. *affinis* type [WFK].

This specimen has lost its BM register number, so that it is not possible to check the accuracy of the locality label 'Barang', which is in Indochina. It is probably wrongly labelled, since the taxon is currently placed as a synonym of *Erythrodiplax funerea* Hagen, a S. American species.

alcestis Tillyard (*Rhyothemis*), 1906 : 482-483, pl. 44, fig. 1. Holotype ♀ (lacking head). Kuranda, N. Q., *F. P. Dodd*, xii.07/*Rhyothemis alcestis* Till., ♀ TYPE, R.J.T.

Described from a single example. Currently placed in the synonymy of *Rhyothemis braganza* Karsch.

allogenes Tillyard (*Agrionoptera*), 1908 : 641-643. LECTOTYPE ♂. Cairns, N. Q., iv.05, *E. Allen*/*Agrionoptera allogenes* Till., TYPE, R.J.T./*Agrionoptera allogenes* Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a subspecies of *A. papuensis* Selys.

amaryllis Selys (*Rhyothemis*), 1878 : 299. Holotype ♂ (lacking head). Menado/Type [McL. label]/Lib. *amaryllis* ♂ Selys.

The unique holotype was originally in the McLachlan collection. Currently placed as a synonym of *Rhyothemis phyllis snelleni* Selys.

amazonica Kirby (*Ephidatia*), 1889 : 331 (pars). LECTOTYPE ♂. Santarem/168/*Ephidatia amazonica* type [WFK]/*Ephidatia amazonica* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

This species was restricted by Ris, 1913 : 1013 to the ♂ and ♀ from Santarem, the ♂ from Para becoming a paratype of *Ephidatia batesi* Ris, 1913. *E. amazonica* Kirby (restr.) is currently considered a synonym of *Idiataphe longipes cubensis* (Scudder).

amazonica Kirby (*Fylgia*), 1889 : 344, pl. 51, figs. 2, 3. LECTOTYPE ♂. Para [label replaced, DEK]/*Fylgia amazonica* type ♂ [WFK]/*Fylgia amazonica* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Kirby's label 'Fylgia amazonica type ♀' had been wrongly attached to an immature ♂ and has been transferred to one of the two ♀ syntypes, both of which lack heads.

ambiguus Kirby (*Misthotus*), 1905a : 193. Lectotype ♂ (Kimmins, 1957 : 96-97). C[ape of] G[ood] H[ope]/*Misthotus ambiguus* cotype [WFK]/*Misthotus ambiguus* Kirby, ♂ Lectotype, D. E. Kimmins det. 1956.

Currently placed as synonym of *Trithemis dorsalis* (Rambur). The other syntype of *ambiguus* (Transvaal) was placed by Longfield (1936 : 490) as a synonym of *Trithemis risi* Longfield (1936 : 490).

anomala Kirby (*Anatya*), 1889 : 338, pls. 53, fig. 9 and 57, fig. 7. LECTOTYPE ♂. Para/3/ Anatya anomala type [WFK]/Anatya anomala Kirby, Lectotype, D. E. Kimmins det. 1967. Currently placed as a synonym of *Anatya guttata* (Erichson).

anomala Pinhey (*Trithemis*), 1955 : 38–39, fig. 8. Holotype ♂. Lake Chila, Abercorn, N. R., iv. 1954, *E. Pinhey*/Trithemis anomala Pinh. 1954, Holotype/Male and Female in Copula.

apicalis Fraser (ssp. *Hylaeothemis fruhstorferi*), 1924. Kimmins, 1966 : 178–179.

apicalis Kirby (*Rhyothemis*), 1889 : 319, pl. 51, fig. 5. Holotype ♀. Aneit[yum], Wallace/apicalis type [WFK].

This taxon is currently placed as a subspecies of *Rhyothemis phyllis* (Sulzer).

apicalis Kirby (*Untamo*), 1889 : 331, pl. 53, fig. 4. Holotype ♀. Sul[a]/Sula, Wallace/Untamo apicalis type [WFK].

Currently placed as a synonym of *Neurothemis ramburi* (Brauer).

ardens McLachlan (*Thecadiplax*), 1894 : 429–430. LECTOTYPE ♂. Ta-chien-lu/Type [McL. label]/Thecadiplax ardens McL./Thecadiplax ardens McL., ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a subspecies of *Sympetrum eroticum* (Selys).

armstrongi Fraser (ssp. *Rhyothemis regia*), 1956. Kimmins, 1966 : 179.

arsinoe Lieftinck (*Diplacina*), 1953 : 171–173, fig. 6. Holotype ♂. Papua, Kokoda, 1,200 ft., viii. 1933, *L. E. Cheesman*/Diplacina arsinoe sp. n. Holotype, det. M. A. Lieftinck, 1947.

atra Pinhey (*Trithemis*), 1961 : 166–167, pl. 11, fig. 14. Holotype ♂. Uganda, Masindi Port, 25. iv. 1956, *P. S. Corbet*/Trithemis atra Pinh., 1956, ♂ Type.

attenuata Kirby (*Trithemis?*), 1889 : 328, pl. 53, fig. 2. LECTOTYPE ♂. Amaz[on]/Trithemis attenuata type ♂ [WFK]/Trithemis? attenuata Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Kirby gives as locality 'Santarem', but the locality labels on the specimens marked by him as ♂ and ♀ types are 'Amaz[on]', and one of the paratypes is labelled as 'Braz.'. In this case I am not interpreting the locality strictly as given in the description but have selected the ♂ labelled by Kirby as type ♂ to be the lectotype. The taxon is currently placed in the genus *Erythrodiplox*.

aurea Fraser (*Tetrathemis*), 1924. Kimmins, 1966 : 180.

auricolor Fraser (var. of *Notiothemis jonesi*), 1944. Kimmins, 1966 : 181.

austeni Kirby (*Perithemis*), 1897 : 602, pl. 12, figs. 4, 5. LECTOTYPE ♂. Manaos, 11. ii. 96, ♂/Manaos, Brazil, 11. ii. 96, *E. E. Austen*/austeni [WFK]/Perithemis austeni Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

austeni Kirby (*Thermorthemis*), 1900 : 72–73, pl. 2, figs. 1, 1a. LECTOTYPE ♂. W. Afr[ica]/Thermorthemis austeni ♂ [WFK].

Kirby based his description upon five males, four from W. Africa (without other locality) and one male, one female from Sierra Leone. Three of the W. African males have been traced and also the male and female from Sierra Leone. This taxon is currently placed as *Orthetrum austeni* (Kirby).

australis Kirby (*Brachydiplax*), 1894 : 18–19. Holotype ♂. Queensland, *Turner*/australis Kb. type [WFK].

australis Kirby (*Brachymesia*), 1889 : 330. Holotype ♂. 'Sydney, N.S.W./australis type [WFK].

Locality label believed to be incorrect and the taxon is currently placed in the synonymy of *Brachymesia furcata* Hagen, a South American species.

azorensis Gardner (ssp. *Sympetrum fonscolombei*), 1958 : 791–792. Holotype ♂. Azores, Pico 13, 17. viii. 52, paired, leading, *J. D. Carthy*/Sympetrum fonscolombei azorensis Gardner, Type ♂, E. Gardner det.

The holotype ♂ and allotype ♀ were taken paired.

azteca Calvert (*Tauriphila*), 1906 : 297, 298. Holotype ♂ (somewhat teneral). Guadalupe, Jalisco, July, *Schumann/Tauriphila azteca* Calv. TYPE. P. P. Calvert det. 1906. B.C.A. Neur., p. 298.

basalis Kirby (*Micrathyria*), 1897 : 610-612, pl. 12, fig. 1. LECTOTYPE ♂. W. end of Parana de Buyassu, 16.i.96/Parana de Buyassu, Lower Amazon, 16.i.96, *E. E. Austen/basalis* [WFK]/*Micrathyria basalis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Kirby mentions two other localities after his description, Obidos (♂) and Breves (♀) and I consider these three examples to be the syntype series. Three additional males discussed later by Kirby are considered as additional non-syntype material, although one of them bears a Kirby cotype label. The taxon is currently placed in the genus *Erythrodiplax*.

basilinea McLachlan (*Libellula*), 1894 : 430-431. LECTOTYPE ♂. Ta-chien-lu/Type [McL. label]/*Libellula basilinea* McL./*Libellula basilinea* McL., ♂ Lectotype, D. E. Kimmins det. 1967.

batesi Kirby (*Cannacia*), 1889 : 341, pls. 53, fig. 1 and 57, fig. 9. Holotype ♂. Amaz[ons]/*Cannacia Batesii* type [WFK].

The ending of the specific name has been changed to a single 'i' in accordance with Art. 31. This taxon is currently placed in the genus *Brachymesia*.

bella Kirby (*Perithemis*), 1889 : 324, pl. 51, figs. 8, 9. LECTOTYPE ♂. 115/Santarem/bella Kb. ♂ [WFK]/*Perithemis bella* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The example in the BM type collection, though bearing Kirby's determination label, was not chosen as lectotype since its locality label is 'Amaz.', whereas the type-locality is specified as 'Santarem'.

bifida Fraser (*Tetrathemis*), 1941. Kimmins, 1966 : 182.

bimacula Kimmins (*Misagria*), 1943 : 156-159, figs. 1-6. Holotype ♂. R. Demarara, B. Guiana/*Misagria bimacula* Kim., ♂ TYPE, det. D. E. Kimmins.

[**biolleyi** Calvert (*Orthemis*), 1906 : 233, 237, pl. 9, figs. 36, 37.

The example in BM(NH) is one of the syntypes but is not labelled Type by Calvert. In the circumstances, Ris (1910 : 286-287) should be considered as having selected the ♂ in the Calvert Collection (Costa Rica, Esparta) as Lectotype, and the ♂ in BM(NH) from Bugaba is therefore a paralectotype.]

blackburni McLachlan (*Lepthemis*), 1882 : 229-231. LECTOTYPE ♂. Hawaiian Islands/*Lepthemis? Blackburni* McL./Type [McL. label]/*Lepthemis blackburni* McLachlan, ♂ Lectotype, D. E. Kimmins det. 1967.

Now placed in the genus *Nesogonia*.

borneense Kimmins (*Orthetrum*), 1936a : 70-73, figs. 2, 3. Holotype ♂. Sarawak, Mt. Dulit, 4,000 ft., 22.x.1932. B in cop. A/Oxford Univ. Exped., *B. M. Hobby, A. W. Moore & J. Ford/Orthetrum borneense* ♂ sp. n., Holotype, det. D. E. Kimmins.

braueri Kirby (*Lyriothemis*), 1889 : 332. Holotype ♀. Sula, Wallace, D. E. Kimmins, 1967/W. Borneo [WFK]/L. Braueri type [WFK].

Kirby gives the locality as Sula [Moluccas] and this locality is repeated by Ris (1909 : 108, 111), after his study of the type. When examining this type during the preparation of this paper, it was found to be labelled 'W. Borneo' in Kirby's writing. It is very unlikely that Ris would not have remarked on the discrepancy in the locality label when he studied the type, and one can only assume that at some later date there was a change of label. I have therefore placed a 'Sula' label above the W. Borneo label. This taxon is currently placed as a synonym of *Lyriothemis cleis* Brauer.

brevistylum Kirby (*Orthetrum*), 1896 : 521-522. Holotype ♂. Dobar, Goolis Mts., 4.ii.1895, *L. L[ort-] P[hillips]/Orthetrum(?) brevistylum* n.s. [WFK].

Currently placed as a synonym of *Orthetrum taeniolatum* (Schneider).

broadwayi Kirby (*Dythemis*), 1894a : 227-228. LECTOTYPE ♂. Trinidad/broadwayi/*Dythemis broadwayi* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
Currently placed as a synonym of *Dythemis velox* Hagen.

burmeisteri Kirby (*Tramea*), 1889 : 316. LECTOTYPE ♂. N.W. India/Burmeisteri Kb. type [WFK]/*Tramea burmeisteri* Kirby, ♂ Lectotype, det. D. E. Kimmins, 1967.

In addition to the allotype ♀, there are 5 paralectotypes in the BM(NH). The taxon is currently placed as a subspecies of *Trapezostigma basilaris* (Palisot de Beauvois).

camarensis Kirby (*Orthetrum*), 1889 : 298. Holotype ♂. Cameroons/camarensis type [WFK]/*Orthetrum camarensis* Kirby, ♂ Holotype, D. E. Kimmins det. 1967.

cambridgei Kirby (*Micrathyria*), 1897 : 608-609, pl. 13, fig. 4. Holotype ♂. Breves, 13.i.96/Breves, Lower Amazon, 13.i.96, *E. E. Austen*/cambridgei [WFK].

campioni Ris (*Allorhizucha*), 1915a : 214-216; Kimmins, 1942 : 47 [selection of holotype]. Lectotype ♂. KaYima, Sierra Leone, 24.vi.1912, *Jas. J. Simpson*/Allorhizucha nova spec. C, Det. Dr. F. Ris/Allorhizucha campioni Ris, ♂ Type, det. D. E. Kimmins.

cannacrioides Calvert (*Dythemis*), 1906 : 272, 276, pl. 8, figs. 43, 44. Holotype ♂. San Isidro, 1,600 ft., *Champion*/Dythemis cannacrioides Calv. ♂ TYPE. P. P. Calvert det. 1906. B. C. A. Neur., p. 277. Orig. Pl. viii, figs. 43, 44.

The penis of the lectotype was removed and studied by Dr. K. Buchholz.

carpenteri Fraser (*Oxythemis*), 1944. Kimmins, 1966 : 186.

carpenteri Fraser (*Tetrathemis*), 1941. Kimmins, 1966 : 186.

celebensis Kirby (*Protorthemis*), 1889 : 334-335, pls. 54, fig. 7 and 57, figs. 6, 6a. Holotype ♂. Mak[assar]*/Celeb[es], Wallace/O. celebensis type [WFK].

ceylonica Kirby (*Zygonidia*), 1905 : 273-275. Holotype ♂. Ceylon. *E. E. Green*/Kandy, ii.1901 [E. E. Green's writing]/Zygonidia ceylonica Kirby, ♂, D. E. Kimmins det. 1961.

Currently placed as subspecies of *Zygonyx iris* Selys.

chirinda Longfield (ssp. *Atoconeura biordinata*), 1953 : 46, pl. 1, figs. 3, B, H. Holotype ♂. Chirinda For., Melsetter Dist., Dept. Agric. S. Rhodesia, 27.xii.1948, *J. A. Whellan*/Atoconeura biordinata chirinda, Type ♂ subsp. nov., det. Miss C. Longfield.

chloe Kirby (*Rhyothemis*), 1894 : 15-16. LECTOTYPE ♀. Queensland, *Turner*/chloe Kirby type [WFK]/Rhyothemis chloe Kirby, ♀ Lectotype, D. E. Kimmins det. 1967.

The lectotype ♀ is the darker of the two syntypes, with (in the hind wing) the 'dark spot on the subnodal sector, halfway between the dark blotch on the nodus and the clouded tip of the wings' mentioned by Kirby. Currently placed as a subspecies of *Rhyothemis phyllis*.

chrysobaphes Ris (*Pseudomacromia*), 1915a : 221-223. LECTOTYPE ♂. Sierra Leone, Sandea, 14.vi.1912, *J. J. Simpson*/Pseudomacromia nova spec. d, Type ♂, Det. Dr. F. Ris/Pseudomacromia chrysobaphes Ris, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed in the genus *Zygonyx*.

cingulata Kirby (*Nesoxenia*), 1889 : 336, pl. 53, fig. 8. LECTOTYPE ♀. [Solomon Islands], Alu/Nesoxenia cingulata type [WFK]/Nesoxenia cingulata Kirby, ♀ Lectotype, D. E. Kimmins det. 1967.

The syntype series also contained 1 ♂ (incomplete) and 1 ♀. This taxon is currently considered a subspecies of *Nesoxenia mysis* Selys.

cladophila Tillyard (*Tetrathemis*), 1908 : 647-648, pl. 14, fig. 5. LECTOTYPE ♂. Cooktown, N. Q., i.08, *R. J. Tillyard*/Tetrathemis cladophila Till. ♂ TYPE. R. J. T./Tetrathemis cladophila Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

* This label was transcribed by Kirby as Makian, but according to Bradley & Betrem, 1967, *Bull. Brit. Mus. nat. Hist. (Ent.)* 20 (7) : 292, the locality should be Makassar.

corduliformis Longfield (*Tetrathemis*), 1936 : 484, figs. 6, A-D. Holotype ♂. Central Africa, Uganda, Bunyoro Distr., Nyamagita Dam, 18.ii.1934, C. E. Longfield/*Tetrathemis corduliformis* sp. n. Type ♂, det. Miss C. Longfield.

coryndoni Fraser (*Lokia*), 1952. Kimmins, 1966 : 187.

cuprina Kirby (*Rhyothemis*), 1889 : 320, pl. 51, fig. 6. Holotype ♀. S. Leone/cuprina Kb. type [WFK].

Currently placed as a synonym of *Rhyothemis fenestrina* (Rambur).

dalei Tillyard (*Nannodythemis*), 1908a : 449-450, pl. 6, fig. 2. LECTOTYPE ♂. Blue Mts., N.S.W., ii.1908, R. J. Tillyard/*Nannodythemis Dalei* Till. ♂ TYPE, R. J. T./*Nannodythemis dalei* Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed in the genus *Nannophya*.

darwinii Kirby (*Tramea*), 1889 : 315, pl. 51, fig. 1. LECTOTYPE ♀. Galapagos/Galapagos/28/Darwinii Kb. type [WFK].

The remaining four female syntypes (in bad condition) are still in the B.M. collection. This taxon is currently placed as a synonym of *Trapezostigma cophysa* (Selys).

davina Fraser (ssp. *Zygonyx iris*), 1926. Kimmins, 1966 : 188.

denticauda Fraser (*Tetrathemis*), 1954. Kimmins, 1966 : 189.

diamangae Longfield (*Aethiothemis*), 1959 : 35, 36, fig. 5A. Holotype ♂ (teneral), N. Angola. Lunda Prov., Dundo, iii.1949/*Aethiothemis diamangae*, Type ♂, det. Miss C. Longfield.

dispar Selys (*Trithemis*, ?aberration), 1883 : 107. Holotype ♀. Japan/Type [McL. label]/*Trithemis phaon*, var. ♀ *dispar* Selys.

Currently placed as a synonym of *Deielia phaon* (Selys).

disparilis Kirby (*Neurothemis*), 1889 : 322-323, pl. 54, fig. 8. LECTOTYPE ♂ (head missing). W. Borneo/N. *disparilis* ♂ [WFK]/*Neurothemis disparilis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

dissocians Calvert (*Micrathyria*), 1906 : 222, 226, pl. 9, figs. 19-21. LECTOTYPE ♂. Atoyac, Vera Cruz, May, H.H.S[mith]/*Micrathyria dissocians* Calv. ♂ TYPE. P. P. Calvert det. 1906. B.C.A. Neur., p. 226, Orig. of pl. ix, figs. 19-21/*Micrathyria dissocians* Calvert, ♂ Lectotype, D. E. Kimmins det. 1967.

Ris (1911 : 440) refers to the Portorico specimen as 'Calvert's Type'. I do not consider this to be a designation of Lectotype, and consider that the data on the above mentioned example makes it clear that Calvert considered it to be the type, especially as the second Vera Cruz example is not marked TYPE.

distanti Kirby (*Stoechia*), 1898 : 236-237. LECTOTYPE ♂. Pretoria (W. L. D[istant])/*distanti* Kirby type [Distant's writing]/*Stoechia distanti* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Trithemis dorsalis* (Rambur).

donaldi Lieftinck (*Lanthanusa*), 1955 : 161-164, figs. 5-9. Holotype ♂. N.E. New Guinea, Saiko, 5,500-6,000 ft., Babu River (Upper Waria River), ix-x.1936, D. Shaw-Meyer/*Lanthanusa donaldi* Lieft. Det. M. A. Lieftinck, 1955. Holotype.

dorothea Fraser (ssp. *Agrionoptera insignis*), 1927. Kimmins, 1966 : 189.

dubia Fraser (*Porpacithemis*), 1954. Kimmins, 1966 : 190.

ducalis Kirby (*Rhyothemis*), 1898 : 230-232. LECTOTYPE ♂. Fort Johnston, Nyassaland, P. Rendall/*Rhyothemis ducalis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Two ♂, one ♀ of the four syntypes are from Fort Johnston; the ♀ from Pretoria, though marked 'cotype' by Kirby, was in effect excluded from the syntypes by Kirby's statement (231) 'The specimens from Fort Johnston are to be considered typical.' Currently placed as a synonym of *R. semihyalina* Desjardins.

eltoni Fraser (*Phyllothemis*), 1935. Kimmins, 1966 : 191.

eludens Tillyard (*Nannophlebia*), 1908 : 645-647, pl. 14, fig. 4. LECTOTYPE ♂. Cairns, N. Q., v.05, *E. Allen*/Nannophlebia eludens Till. ♂ TYPE, R.J.T./Nannophlebia eludens Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

equestris Fabricius (*Libellula*), 1781 : 523. *Campion*, 1917 : 445.

equivocata Kirby (*Thermochoria*), 1889 : 339, pl. 52, fig. 8. Holotype ♂. W. Afr[ica]/Thermochoria equivocata type [WFK].

erichsoni Kirby (*Trithemis*), 1894 : 263. LECTOTYPE ♂. Amaz./erichsoni [WFK]/Trithemis erichsoni Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

This specimen was the basis of the drawings and notes sent by the present author to D. J. Borror in Jan. 1935, and referred to in his monograph on *Erythrodiplax*. This taxon is currently placed in synonymy of *Erythrodiplax unimaculata* (De Geer).

eudoxia Kirby (*Accaphila*), 1909 : 60. Holotype ♂. Ruwenzori East, alt. 6,000 ft., 30.i.07/Ruwenzori Exp./Accaphila eudoxia Kb. type.

Currently placed in the genus *Atoconeura*.

eurybia Selys (*Libellula*), 1878 : 293, 298. Holotype ♂. Menado/Type [McL. label]/Lib. eurybia Selys ♂, race de Chinensis? [Selys writing].

Currently placed in the genus *Trapezostigma*.

exigua Kirby (*Fylla*), 1889 : 345, pl. 52, fig. 6. LECTOTYPE ♂. Gil[olo]/Wallace/Fylla exigua Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The syntype series was found in the museum collection, without type-indication or determination labels by Kirby. One can only presume that as Kirby reduced the taxon to synonymy of *Nannophya pygmaea* Rambur in his Catalogue of Odonata published the following year, he removed his determination labels. The series now contains paralectotypes from Borneo, Gilolo and Morty.

eximia Kirby (*Micrathyria*), 1897 : 609, pl. 13, fig. 3. LECTOTYPE ♂. Obydos, 2.ii.96/Obydos, Lower Amazon, 2.ii.96, *E. E. Austen*/eximia [WFK]/Micrathyria eximia Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The lectotype is the specimen which, for many years, has carried a red BM type label. The remaining 3 ♂ syntypes have been labelled paralectotypes. Since Kirby stated that the species was described from four males from Obydos, the male mentioned by him from Para is excluded from the syntype series.

extraordinata Fraser (*Atoconeura*), 1950. *Kimmins*, 1966 : 191.

falsum Longfield (ssp. *Orthetrum capense*), 1955 : 26. Holotype ♂. B. E. Africa, Kenya, Thoura Forest, Meru, 600 ft., 30.i.1934, far from water, *C. E. Longfield*/Orthetrum capense falsum Longf., Type ♂, det. Miss C. Longfield.

Currently placed as subspecies of *Orthetrum capicola* Kimmins.

fasciata Kirby (*Celithemis*), 1889 : 326, pl. 52, fig. 2. LECTOTYPE ♂. Georgia/fasciata Kb. type [WFK]/Celithemis fasciata Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

fasciata Kirby (*Deielia*), 1892 : 330, pl. 53, fig. 6. McLachlan, 1892 : 177-178. Holotype ♀. 137/Sandw[ich] Is., *Beechey*/Loo Choo Is., Voy. of 'Blossom', See McLachlan, *An. Mag. Nat. Hist.* (6) x. p. 177 (1892)/Deielia fasciata, type [WFK].

As explained by McLachlan (1892), the original locality label on this specimen appears to have been incorrect, as the insect has not been refound in the Hawaiian Islands. It does occur in the Loo Choo (Ryukyu) Islands, which were also visited by the 'Blossom', and it is considered most likely that the example came from these islands. The taxon is currently considered a synonym of *Deielia phaon* (Selys).

ferruginata Fabricius (*Libellula*), 1781 : 521. *Campion*, 1917 : 445.

fitzgeraldi Pinhey in Longfield (*Nesciothemis*), 1955 : 61-63, figs. 10 B, D, F; Pinhey, 1955 : 30-32, fig. 6. Holotype ♂. Lake Chila, Abercorn, N. R., 31.i.54, *D. Vesey-Fitzgerald*/Orthetrum(?) fitzgeraldi Pinh. 1954, Holotype/Nesciothemis fitzgeraldi (Pinhey), det. Miss C. Longfield.

The type of *Orthetrum*(?) *fitzgeraldi* Pinhey was presented to the BM(NH) when the manuscript was sent to press. Publication was unavoidably delayed, and in the belief that Pinhey's paper was due to appear, Miss Longfield included a description of *fitzgeraldi* in her paper on African *Orthetrum*, placing it in her new genus *Nesciothemis*, the publication of which unfortunately antedated Pinhey's paper.

flava Longfield (*Monardithemis*), 1947 : 23-25, figs. 9-10. Holotype ♂ (original designation as type). Kalukembé, Dec./Angola, Miss. sc. suisse, 1932-1933/Monardithemis flava genotype ♂, det. Miss C. Longfield.

flavescens Kirby (*Miathyria*), 1897 : 600-601, pl. 13, fig. 2. LECTOTYPE ♂. Santarem/1113 ♂/Miathyria flavescens Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The lectotype ♂, which has for many years carried a red BM type-label, has no determination label by Kirby, but the locality label agrees with the type-data and there is no doubt that it is one of the syntypes. On the other hand, an example bearing Kirby's label 'Miathyria flavescens K. cotype' is from Para, not Santarem and is from the Saunders collection (not Bates) and cannot therefore be a syntype.

flavescens Kirby (*Tetrathemis*), 1889 : 343, pl. 52, fig. 4. LECTOTYPE ♂. Sar[awak]/Wallace/T. flavescens type [WFK]/Tetrathemis flavescens Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

flavidulum Kirby (*Orthetrum*), 1898 : 238. LECTOTYPE ♀. Pretoria, W. L. D.[istant]/Orthetrum flavidulum Kirby, ♀ Lectotype, D. E. Kimmins det. 1967.

Of the four examples (listed by Kirby as females), the example from the Pienaars River, iii. 1894 proves to be a ♂. Currently placed as synonym of *Orthetrum abbotti* Calvert.

flavifrons Kirby (*Chalcostephia*), 1889 : 337. Holotype ♂. Angola/Chalcostephia flavifrons type.

flavopicta Kirby (*Orthemis*), 1889 : 332-333, pls. 54, fig. 1 and 57, fig. 5. LECTOTYPE ♂. Para/43/Orthemis flavopicta type/Orthemis flavopicta Kirby, Holotypus präp. Dr. Buchholz/Orthemis flavopicta Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

From the syntype series of 4 ♂, 1 ♀, the ♂ bearing Kirby's type label has been chosen as lectotype.

foliata Kirby (*Belonia*), 1889 : 333, pl. 54, fig. 4. Holotype ♀ (by original designation). [Mexico], Dueñas/foliata Kb. type [WFK].

Currently placed in the genus *Libellula*. The ♂ from Guatemala, doubtfully referred to *Belonia foliata* by Kirby, is now considered to be a synonym of *Libellula herculea* Karsch.

fraseri Pinhey (*Olpogastra*), 1955 : 39-40, pl. 3, fig. 1, text-fig. 8. Holotype ♂. Aswa R., Acholi, Uganda, March 1952, T. H. E. Jackson. *Olpogastra fraseri* Pinh. 1954, Holotype.

Currently placed in the genus *Zygonoides* Pinhey.

frontalis Kirby (*Lyriothemis*), 1889 : 332. Holotype ♀. Sul[a]/Sula, Wallace/L. frontalis type [WFK].

Currently placed as a synonym of *Lyriothemis cleis* Brauer.

fulgens Kirby (*Rhyothemis*), 1889 : 322. Holotype ♂. Sar[awak]/fulgens, Kb. type [WFK].

gamblesi Longfield (*Oxythemis*), 1959 : 36-38, figs. 5 B, C. Holotype ♂. S. Nigeria, Ikorodu, Ogun Delta, 26.ii.1956, R. M. Gambles/Oxythemis gamblesi Type ♂, det. Miss C. Longfield.

gardneri Fraser (*Hylaeothemis*), 1927. Kimmins, 1966 : 194.

grenadensis Kirby (*Brechmorhoga*), 1894b : 265. LECTOTYPE ♂. Grenada/106/grenadensis [WFK]/Brechmorhoga grenadensis Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a subspecies of *Brechmorhoga praecox* (Hagen).

[**hageni** Kirby (*Micrathyria*), 1890 : 41]

This name was proposed for *Dythemis didyma* Hagen, 1861, nec Selys, 1857 and the type should therefore be selected from Hagen's material.

hyalinum Kirby (*Orthetrum*), 1886 : 326-327, pl. 33, figs. 5, 6. LECTOTYPE ♂. N.W. Ind[ia]/Campbellpore, 14. xi. 85, 7/hyalinum Kirb. ♂ type [WFK]/*Orthetrum hyalinum* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a synonym of *Orthetrum taeniolatum* (Schneider).

hyalina Kirby (*Tetrathemis*), 1889 : 342, pl. 56, fig. 8. Holotype ♂ (lacking abdominal segments 5-10). Borneo/T. *hyalina* type [WFK].

Currently placed as subspecies of *Tetrathemis irregularis* Brauer.

[*icteroptera* Selys (*Libellula*), 1857, in Sagra, *Hist. Cuba*, 7 : 190.]

This species was briefly differentiated as a race in a note on *Libellula metella* Selys; the type locality being given as Buenos Ayres. The McLachlan collection included an example determined by Selys as *Lib. icteroptera* Selys ♀, to which McLachlan had added a type label. This cannot however be the type, as its locality label is Minas Geraes.

immaculata Fraser (*Zygonyx*), 1933. Kimmins, 1966 : 196.

imperatrix Selys (*Rhyothemis*), 1887 : 53-54. Holotype ♂. Ryukyu [Loo Choo], vi-viii. 1887 [Pryer]/*Rhyothemis imperatrix* Selys, nov. sp.? ou *race de splendida*, R. ♂, Loo Choo [Selys' writing].

Selys saw originally only this one example, but McLachlan subsequently received three females with the same locality data (collected by Pryer), of which one was given to Selys and was described by him (1888 : 1); the other two females are now in BM(NH).

indica Fabricius (*Libellula*), 1781 : 521. Campion, 1917 : 444.

indica Fraser (*Hylaeothemis*), 1946. Kimmins, 1966 : 196.

indica Kirby (*Brachydiplax*), 1889 : 329, pl. 54, fig. 9. LECTOTYPE ♂. N. Ind[ia]/B. indica ♂ type [WFK]/*Brachydiplax indica* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The mature ♂ has been chosen as lectotype, as this example has been segregated in the type-collection since 1940.

infuscatum Selys (*Diplax*), 1883 : 90. LECTOTYPE ♂. Japan/Type/*Diplax infuscata* Selys, ♂ Japon/*Diplax infuscata* Selys, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed in the genus *Sympetrum*.

insignis Kirby (*Zygonidia*), 1900a : 533-534, pl. 12, fig. 1. LECTOTYPE ♂. 5-fingered Mt., Hainan, J. T. Thomasson/*Zygonidia insignis*, type [WFK]/*Zygonidia insignis* Kirby, ♂ Lectotype, D. E. Kimmins det., 1967.

Currently placed as a synonym of *Zygonyx iris* Selys.

insularis Kirby (*Agrionoptera*), 1889 : 336-337. LECTOTYPE ♂. [Solomon Is.] Santa Anna/*Agrionoptera insularis* ♂ type [WFK]/*Agrionoptera insularis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a subspecies of *Agrionoptera insignis* (Rambur).

insularis Kirby (*Cannaphila*), 1889 : 341. Lectotype ♀ (Ris, 1910 : 295). Hayti/*Cannaphila insularis* Kirby, ♀ Lectotype, F. Ris design. 1910 [D.E.K. hand].

I consider that Ris' action (1910 : 295) in citing 'British Museum: 1 ♀ juv. Hayti (Kirby's Type)' is equivalent to selection of a lectotype. There is one other ♀ in BMNH, which bore Kirby's determination *Cannaphila insularis*, but this has a locality label St. Dom. [Santo Domingo], which is in Haiti and was probably a syntype. This is now labelled paralectotype. I have been unable to trace the ♂ syntype referred to by Kirby.

Kirby's statement '*Hab. Haiti* (types in British Museum); Jamaica (Dublin Museum)' eliminated the Dublin Museum example(s) from consideration as lectotype.

intensa Kirby (*Perithemis*), 1889 : 326, pl. 51, fig. 7. Holotype ♂ (abd. segs. 9-10 missing). Mex[ico]/*intensa* Kb. type [WFK].

intermedia Fraser (*Protorthemis*), 1936. Kimmins, 1966 : 198.

internum McLachlan (var. of *Orthetrum japonicum*), 1894 : 431-432. LECTOTYPE ♂. Ta-chien-lu/O. japonicum var. internum McL./Orthetrum japonicum internum McL., ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as *Orthetrum japonicum internum* McL.

iridescens Kirby (*Urothemis*), 1898 : 235. LECTOTYPE ♂ (abd. segs. 6-10 missing). Fort Johnston, Nyassaland, *P. Rendall/iridescens* Kirby type [WFK]/*Urothemis iridescens* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Urothemis edwardsi* (Selys).

isa Fraser (ssp. of *Zygonyx iris*), 1926. Kimmins, 1966 : 198.

isis Fraser (*Zygonyx*), 1924. Kimmins, 1966 : 198.

ulia Kirby (*Orthetrum*), 1900 : 75-77, pl. 2, fig. 3. LECTOTYPE ♂ (abd. segs. 9-10 missing). Sierra Leone/Orthetrum julia type [WFK].

Kirby writes (: 77) 'The species is described from four specimens from Sierra Leone, two of which were collected by Mr. Austen on Aug. 26 and Sept. 11 (1899).' The example designated as Lectotype is the only one bearing Kirby's determination label. The allotype female has for many years borne a museum type label but now has no locality label (?Sierra Leone, 42.31). Of the listed examples collected by Austen, one (without abdomen), labelled 'Free Town, Sierra Leone, 26. viii. 1899, E. E. Austen' has been located but the other is missing.

kalai Longfield (ssp. *Orthetrum stemmale*), 1936 : 487, 488, fig. 7. Holotype ♂. S. Rhodesia, Victoria Falls, Zambesi River, Kalai Island, 7.iv.1936, *C. E. Longfield/Orthetrum stemmale kalai* sp. nov. Type ♂, det. Miss C. Longfield.

kalula Kirby (*Trithemis*), 1900 : 69, pl. 2, figs. 2, 2a. LECTOTYPE ♂. Free Town, Sierra Leone, 13.ix.99, *E. E. Austen/Trithemis kalula* Kb. type [WFK].

Kirby lists four males, but only three can now be traced in the collections; of these, only the lectotype bears Kirby's determination label. This specimen has for many years carried a red type label and has been considered to be the type. The two remaining males are considered paralectotypes.

kenya Longfield (spp. *Atoconeura biordinata*), 1953 : 46, pl. 1, figs. 5, 9, C, G. Holotype ♂. B.E. Africa, Kenya, Meru, 6,000 ft., on swift stream, 29.i.1934, *C. E. Longfield. Atoconeura biordinata kenya*, Type ♂ subsp. n., det. Miss C. Longfield.

kerri Fraser (*Amphithemis*), 1933. Kimmins, 1966 : 198.

lacustris Kirby (*Trithemis?*), 1889 : 329. Holotype ♂. [Central Africa] Wadelai, 27.i.1887 [*Emin Pasha*]/lacustris Kb. type [WFK].

Currently placed in the genus *Brachythemis*.

lankana Kirby (*Rhyothemis*), 1893 : 549-550. Holotype ♂. Udagama, 26.iv.92/Ceylon, Yerbury, (on reverse) lankana type/Rhyothemis lankana Kirby, ♂ Holotype, D. E. Kimmins det. 1967.

leakeyi Pinhey (*Lokithemis*), 1955 : 34-35, pl. 3, figs. 3, 4, 6, and text-fig. 7. Holotype ♂. Lake Chila, Abercorn, N. Rhodesia, 31.i.54, *D. Vesey Fitzgerald/Lokithemis leakeyi* Pinh. Holotype.

[**levis** Calvert (*Orthemis*), 1906 : 233, 238, pl. 9, figs. 38, 39.]

No example in our B.C.A. series was marked TYPE by Calvert and our specimens are therefore considered as syntypes. The true type is probably in some other collection.

lewisi Selys (*Lyriothemis*), 1883 : 96-97. LECTOTYPE ♂. Japan, *Lewis/Lyriothemis lewisii* Selys ♂, Japon/Lyriothemis lewisi Selys, ♂ Lectotype, D. E. Kimmins det. 1967.

Ris (1909 : 119) in the list of material dealt with cites '1 ♂ Japon (Pryer, Selys Type der *L. lewisii*)'; this cannot be accepted as a designation of lectotype, since Selys limits his material to 'Japon, par M. G. Lewis, communiquée par MacLachlan (coll. MacLachlan, Selys)'. I have therefore designated the above specimen as Lectotype. Currently placed as a synonym of *L. pachygastra* (Selys).

- lieftincki** Fraser (*Rhodothemis*), 1954. Kimmins, 1966 : 201.
- lifuana** Kimmins (ssp. *Agrionoptera insignis*), 1953 : 243-244, figs. 1-3. Holotype ♂. Lifu/Agrionoptera insignis lifuana Kim. ♂ Holotype, D. E. Kimmins det. 1951.
Currently placed as a subspecies of *Agrionoptera papuensis*.
- longipennis** Kirby (*Belonia*), 1889 : 334. Holotype ♀. Cuença/longipennis Kb. type [WFK].
Currently placed as a synonym of *Libellula herculea* Karsch.
- lori** Kirby (*Orthetrum*), 1896 : 522-523. Holotype ♀ (lacking abd. segs. 6-10). Dobar, Goolis Mts., 4.ii.1896, L. L[ort-] P[hillips]/Orthetrum lorti.
Currently placed as synonym of *Crocothemis erythraea* (Brullé).
- machadoi** Longfield (*Orthetrum*), 1955 : 35-39, figs. 5, A, E, J. Holotype ♂. N.E. Angola, Dundo, Lunda Distr., 13.xii.1947/Orthetrum machadoi Type ♂, in cop. with type ♀, det. Miss C. Longfield.
- macrostigma** Longfield (*Orthetrum*), 1947 : 25-27, fig. 11. Holotype ♂. Lunda, sept./Angola Miss. sc. suisse, 1932-1933/Orthetrum macrostigma Type ♂, det. Miss C. Longfield.
- madagascariensis** Kirby (*Tramea*), 1889 : 317. Holotype ♂. Betsileo, Madag. [WFK]/madagascariensis Kb. type [WFK].
This taxon is currently placed as a synonym of *Trapezostigma limbata* (Desj.).
- malabarica** Fraser (ssp. *Zygonyx iris*), 1926. Kimmins, 1966 : 203.
- malcolmi** Fraser (*Brachydiplax*), 1922. Kimmins, 1966 : 203.
- marginata** Fabricius (*Libellula*), 1781 : 523. Champion, 1917 : 445.
- marshalli** Kirby (*Misthotus*), 1905a : 192-193. LECTOTYPE ♂. Salisbury, Mashonaland, Marshall/5. Salisbury, Nov. 1903, in copula. *G. A. K. M[arshall]/marshalli* Kb. type [WFK]/Misthotus marshalli Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
The 'still more immature male' referred to by Kirby cannot be traced. Currently placed as a synonym of *Trithemis dorsalis* (Rambur).
- maya** Calvert (*Dythemis*), 1906 : 272, 275, pl. 8, fig. 45. Holotype ♂. San Geronimo, Vera Paz, *Champion/Dythemis maya* Calv. TYPE ♂. P. P. Calvert det. 1906. B. C. A. Neur., p. 275. Original of Pl. viii, fig. 45.
The penis of the lectotype was removed and studied by Dr. Buchholz.
- mallicca** Fraser (ssp. of *Zygonyx iris*), 1931. Kimmins, 1966 : 203.
- mildredae** Fraser (ssp. of *Zygonyx iris*), 1920. Kimmins, 1966 : 204.
- monteiroi** Kirby (*Thermorthemis*), 1900 : 73. Holotype ♀. Angola/monteiroi Kb. type [WFK].
Currently considered a synonym of *Orthetrum austeni* (Kirby).
- multipunctata** Kirby (*Dythemis*), 1894b : 265-266. LECTOTYPE ♂ (left fore wing missing). St. Vincent/Pools near Bannonallie, Jan. 12/multipunctata [WFK]/Dythemis multipunctata Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
I have chosen as lectotype the ♂ which has for many years carried a BM red type label and of which the penis was removed and studied by Dr. K. Buchholz.
- multipunctata** Kirby var. (*Dythemis*), 1894b : 266. LECTOTYPE ♂. Grenada/235/Dythemis multipunctata var. Kirby, ♂ Lectotype, D. E. Kimmins det. 1967/Dythemis sterilis Hag., det. D. E. Kimmins.
Currently placed as a synonym of *Dythemis sterilis* Hagen.
- nicevillei** Kirby (*Orthetrum*), 1894c : 112-113. LECTOTYPE ♂. Burma/nicevillei Kb. type ♂ [WFK]/Orthetrum nicevillei Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
One of Kirby's determination labels had at some time been transferred from a ♂ syntype to a ♀ which was not received at the BM(NH) until after publication of the description. This label has been replaced on its correct specimen. Currently placed as synonym of *Orthetrum glaucum* Brauer.

nigra Longfield (ssp. of *Trithemis donaldsoni*), 1936 : 491-493, 496, figs. 9, 10. Holotype ♂ (last 5 abdominal segments missing). W. Africa, Principe I., Ogui Pipi, 7.xii.1932, *W. H. T. Tams/Trithemis donaldsoni nigra* ssp. nov. Type ♂, det. Miss C. Longfield.

In the original description, only the above example was mentioned, but in a list (: 496), two ♂ are recorded. The second one (a complete example) had been labelled by Miss Longfield as type and the holotype labelled as paratype. This error has now been corrected and the second example cannot be considered even as a paratype, as it is not mentioned in the original description.

nigricolor Fraser (*Amphithemis*), 1922. Kimmins, 1966 : 206.

nigrifrons Kirby (*Orthetrum*), 1894 : 19-20. LECTOTYPE ♂. Queensland, *Turner/Orthetrum nigrifrons* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The lectotype and paralectotype lack any determination label by Kirby. Currently placed in the genus *Crocothemis*.

nigrilabris Selys (*Lib[ellula]*), 1872 : 177; (*Urothemis*), Selys, 1878 : 304-305. Holotype ♀. Menado/Lib. *nigrilabris* Selys, ♀ Type/*Macrodiplax lycoris* Selys.

This taxon was first described briefly (1872) in comparison with *Libellula lycoris* Selys, and a full description, with location of type given, in 1878. Currently placed as synonym of *Macrodiplax cora* Brauer.

notabilis Kirby (*Onychothemis*), 1906 : 277-278. Holotype ♂. Tonkin [*Fruhstorfer/Onychothemis notabilis* Kirby, Holotype ♂, det. D. E. Kimmins, 1967.

No Kirby determination label. Currently placed as synonym of *Onychothemis tonkinensis* Martin, 1904.

notata Fabricius (*Libellula*), 1787 : 337. Campion, 1917 : 444.

novaezealandiae McLachlan (var. of *Sympetrum bipunctatum*), 1894a : 271-272. LECTOTYPE ♀. 29e[Hudson's locality number]/Paikakariki, Wellington, N.Z., *Hudson/Sympetrum bipunctatum* Br., var. *novae-zealandiae* McL. Type/*Sympetrum bipunctatum* var. *novaezealandiae* McL., ♀ Lectotype, D. E. Kimmins det. 1967.

The fourth syntype (from Auckland) has not been traced. The name *novaezealandiae* was originally hyphenated and has been emended in accordance with Art. 32(e)(i). Currently placed as a synonym of *Diplacodes bipunctata* (Brauer).

obsolescens Kirby (*Rhyothemis*), 1889 : 321. Holotype ♂. Borneo/*obsolescens* type [WFK].

occidentalis Tillyard (*Nannodythemis*), 1908a : 450-453, pl. 6, fig. 3. LECTOTYPE ♂. Bridgetown, W. A[ustralia], i.07, *R. J. Tillyard/Nannodythemis occidentalis* Till., ♂ TYPE. R.J.T./*Nannodythemis occidentalis* Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed in the genus *Nannophya*.

octomaculata Fraser (*Palpopleura*), 1935. Kimmins, 1966 : 207.

oculata Fabricius (*Libellula*), 1775 : 421. Campion, 1917 : 444.

osiris Fraser (ssp. *Zygonyx iris*), 1936. Kimmins, 1966 : 208.

othello Tillyard (*Camacinia*), 1908 : 639-641, pl. 14, fig. 1. Holotype ♂. Cooktown, N.Q., xii.07, *R. J. Tillyard/Camacinia Othello* Till. ♂ TYPE, R. J. T.

pacificus Kirby (*Diplax*), 1884 : 355. Holotype ♂. Tongatabu, with on reverse, *Diplax pacificus* type [WFK].

Currently placed as synonym of *Diplacodes bipunctatus* (Brauer).

pallidinervis Kirby (*Sympetrum*), 1889 : 327, pl. 55, fig. 4. LECTOTYPE ♂. Ind[ia]/*Pseudothemis pallidinervis* Kb. type [WFK]/*Sympetrum pallidinervis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as *Trithemis pallidinervis* (Kirby). Although described as a *Sympetrum*, Kirby labelled his specimens as *Pseudothemis*.

parana Kirby (*Misagria*), 1889 : 339, pls. 52, fig. 9 and 57, fig. 8. Holotype ♂. Para/Misagria parana type [WFK].

parasticta Pinhey (*Trithemis*), 1955 : 35-37, fig. 8. Holotype ♂. Lake Chila, Abercorn, N. R., 31. i. 54, *D. Vesey Fitzgerald/Trithemis parasticta* Pinh., 1954.

phaon Selys (*Trithemis*), 1883 : 106-107. LECTOTYPE ♂. Yokohama, Orio/Type [McL. label]/*Trithemis phaon* Selys ♂, Japon [Selys label]/*Trithemis phaon* Selys, ♂ Lectotype, D. E. Kimmins det. 1967.

Selys gives no indication as to the collection in which the type was deposited. Ris (1911 : 589) makes no mention of the type in the Selys Collection and therefore the above specimen from the McLachlan Collection has been designated as Lectotype. Currently placed in the genus *Deielia* Kirby.

phillipsi Kirby (*Orthetrum*), 1896 : 522. Holotype ♂. Dobar, 2nd Feby. 95, maritime plain/Dobar, Goolis Mts, 2. ii. 1895, *L. L[ort-] P[hillips]/Orthetrum Phillipsi* Kb. type [WFK].

It may be pointed out that the locality data given on the second (printed) label and in the description differs from that given on the first (mss) label. The date on the printed label has been altered to conform with the mss label. Currently placed as a synonym of *Orthetrum abbotti* (Calvert).

prateri Fraser (*Pseudotrimea*), 1920. Kimmins, 1966 : 210.

princeps Kirby (*Rhyothemis*), 1894 : 16-17. LECTOTYPE ♂. Queensland, *Turner, princeps* Kb. ♂ type/*Rhyothemis princeps* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

pruinosa Kirby (*Micrathyria*?), 1894b : 267-268. Holotype ♂ (abd. segments 8-10 missing). Grenada/233/pruinosa [WFK]/*Micrathyria* (?) *pruinosa* Kirby, ♂ Holotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Micrathyria didyma didyma* (Selys).

pseudeudoxia Longfield (ssp. *Atoconeura biordinata*), 1953 : 46, pl. 1, figs. 2, E, H. Holotype ♂. Uganda, Ruwenzori Range, Mubuku Valley, Bikoni Peak, 7,000 ft., 30. xii. 1937, *C. E. Longfield/Atoconeura biordinata pseudeudoxia* Type ♂, ssp. nov. det. Miss C. Longfield.

pseudodeflecta Pinhey (ssp. *Hadrothemis deflecta*), 1961 : 127-128, pl. 9, fig. 2. Holotype ♂. Entebbe, Uganda, Oct. 1952, *E. Pinhey/Hadrothemis deflecta pseudodeflecta* Pinh. Holotype.

puella Kirby (*Aino*), 1890 : 113. LECTOTYPE ♂ (lacking abd. segs. 9-10). *N. pygmaea/Aino puella* Kirby, ♂ Lectotype. D. E. Kimmins det. 1967.

The only example labelled 'puella' and with type label was the female, but the ♂ syntype was discovered labelled solely 'N. pygmaea'. It agrees well with the original description and has been designated the lectotype. The locality of the syntypes was unknown. The taxon is currently placed as a synonym of *Nannothemis bella* Uhler, a South American insect.

pusilla Kirby (*Miathyria*), 1889 : 318, pl. 52, fig. 3. LECTOTYPE ♂. Amaz[ons]/142/*Miathyria pusilla* type [WFK]/*Miathyria pusilla* Kirby, ♂ Lectotype, det. D. E. Kimmins, 1967.

Kirby gives as localities for this species, Tapajos and Santarem; there are three paralectotypes from these localities, one of which is labelled 142 ♀. In spite of the fact that the specimen bearing Kirby's type label has only 'Amaz[ons]' as locality, I have accepted it as lectotype. This taxon is currently placed as a synonym of *Miathyria simplex* (Rambur).

reducta Fraser (*Uracis*), 1946. Kimmins, 1966 : 211.

regalis Tillyard (*Agrionoptera*), 1908 : 643-645, pl. 14, fig. 3. LECTOTYPE ♂. Cooktown, N.Q., i. 08, *R. J. Tillyard/Agrionoptera regalis* Till., ♂ TYPE, R.J.T./*Agrionoptera regalis* Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Agrionoptera longitudinalis* Selys.

rendalli Kirby (*Urothemis*), 1898 : 234. Holotype ♀. Fort Johnston, Nyassaland, *P. Rendall/rendalli* Kirb. paratype [Distant's writing]/*Urothemis rendalli* Kirby ♀ TYPE, D. E. Kimmins det. 1956.

As *U. rendalli* was described from one female, Distant's label 'paratype' must be an error. Currently placed as synonym of *Urothemis edwardsi* (Selys).

[*resplendens* Selys (*Rhyothemis*), 1878 : 300–301.]

The unnamed variety mentioned by Selys (: 301) is in the BM(NH). Although given a type label by McLachlan, it is not considered to be a type.

reticulata Kirby (*Crocothemis*), 1886 : 328, pl. 33, figs. 8, 9. Holotype ♂. Campbellpore, 21.xi.85/N.W. Ind[ia]/reticulata Kirb. type.

Currently placed as synonym of *Crocothemis servilia* (Drury).

rezia Kirby (*Aethriamanta*), 1889a : 298. Holotype ♂ (lacking abd. segs. 5–10). Madag[ascar]/rezia Kb. type.

risi Champion (*Oda*), 1915 : 489–491, fig. 40. Holotype ♂. Utakwa R., Dutch N. Guin., 2,500–3,000 ft., ii.1913, *A. F. R. Wollaston*/Oda nov. spec. Det. Dr. F. Ris/Oda risi Cmpn. Holotype, Determined by H. Champion.

Currently placed in the genus *Risiophlebia*.

risi Longfield (*Trithemis*), 1936 : 490, 491, fig. 8A. Holotype ♂. Brit. E. Africa, 31.iii.11, *S. A. Neave*/Trithemis distantis ♂, Dr. F. Ris det./Trithemis risi Longfield, ♂ Holotype, D. E. Kimmins det. 1967.

Longfield (: 491) states 'Lectotype ♂ and ♀ have been selected from specimens collected by Dr. Neave in British E. Africa in 1911.' As she was describing a new species, in effect she designated holotype ♂ and allotype ♀. Pinhey (1962, *Publ çôes cult. Co. Diam. Angola* 59 : 271) quotes the Transvaal ♂ of *ambiguus* Kirby as lectotype of *risi* Longfield, but although Longfield quotes this specimen in synonymy, she designated a holotype for *risi* Longfield.

risi Pinhey (ssp. *Porpax asperipes*), 1958a : 115–116. Holotype ♂. Chibudzana R., Melsetter Dist., 26.ii.1948, Dept. Agric. S. Rhodesia/Porpax asperipes ssp. risi Pinhey, 1956, Type.

Subsequently raised to specific rank.

risi Tillyard (*Nannophlebia*), 1913a : 713–714, pl. 74, figs. 4, 5, 7. LECTOTYPE ♂. Belling R., N.S.W., 30.xi.11, *R. J. Tillyard*/Nannophlebia risi Till., ♂ TYPE, R. J. T./Nannophlebia risi Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.

robertsi Fraser (*Notiothemis*), 1944. Kimmins, 1966 : 211.

rubra Kirby (*Trithemis*), 1889 : 328. LECTOTYPE ♂. Australia/T. rubra type [WFK]/Trithemis rubra Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

As Australia is the first locality mentioned, and Kirby's type label was attached to the Australian specimen, this has been chosen as the Lectotype. Currently placed as a synonym of *Diplacodes haematodes* Burmeister.

ruwenzoriensis Fraser (*Tetrathemis*), 1941. Kimmins, 1966 : 211.

sagitta Ris (*Orthetrum*), 1915a : 216–217. LECTOTYPE ♂. Sierra Leone, Port Lokko, 3.v.1912, *J. J. Simpson*/Orthetrum nov. spec. ♂ (Type) (near africanum)[Ris' writing]/Orthetrum sagitta Ris, ♂ Lectotype, D. E. Kimmins det. 1967.

The penis of the lectotype was dissected and studied by Dr. K. Buchholz, 1963.

schumanni Calvert (*Micrathyria*), 1906 : 225, 227, pl. 8, figs. 39, 40, pl. 9, fig. 24. Holotype ♂. Guadalajara, Jalisco, July, *Schumann*/Micrathyria schumanni Calv., TYPE. P. P. Calvert det. 1906. *B. C. A. Neur.*, p. 227, Orig. pl. viii, ff. 39, 40, pl. ix, fig. 24.

The ♂ right hamule has been removed and mounted in canada balsam.

serva Kirby (*Trithemis*), 1900 : 69–70, fig. 1. LECTOTYPE ♂ (abd. segs 7–10 missing). Free Town, Sierra Leone, 5.ix.99, *E. E. Austen*/On reverse, T. serva K[WFK].

Kirby is a little obscure in his account of the material upon which this taxon is based. He says 'Described from six specimens, two of them taken by Mr. Austen on Sept. 5 and 21 (1899).' These two specimens are in the collections; one is labelled 'Trithemis serva Cotype' in Kirby's writing, the other (with a red type label) has 'T. serva K' in Kirby's writing on underside of locality label. There is also another Sierra Leone specimen, collected by Morgan, 1835, labelled 'Trith. serva' by Kirby. Three other Free Town examples, collected by Austen, bear no determination labels and I consider them to be excluded from the type series by Kirby's

statement that *two* were taken by Austen. One must assume that the other three examples are missing. I have therefore chosen the specimen which originally bore the red type label as Lectotype.

siamensis Fraser (ssp. of *Onychothemis tonkinensis*), 1932. Kimmins, 1966 : 213.

siemensi Kirby (*Uracis*), 1897 : 605-606, pl. 12, fig. 2. LECTOTYPE ♂. Gurupa, Lower Amazon, 22.ii.96, *E. E. Austen/siemensi* [WFK]/*Uracis siemensi* holotypus, präp. Dr. Buchholz/*Uracis siemensi* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The penis of the lectotype was removed and studied by Dr. H. Buchholz on one of his visits to BM(NH). The 'female from Para' referred to by Kirby is now labelled allotype and the second male (marked cotype by Kirby) has been labelled paralectotype.

simpsoni Ris (*Cyanothemis*), 1915 : 219-221, fig. Holotype ♂ (original designation). Yana, Sierra Leone, 1.iv.1912, *Jas. J. Simpson/Nov. gen. nov. spec.c.*, Det. Dr. F. Ris/*Cyanothemis simpsoni* Ris, ♂ Holotype, D. E. Kimmins det. 1967.

Ris (1919 : 219) refers to this Yana specimen in his description of the subjuvenile ♂ from Mongheri as 'Like first(type) specimen,' and also figures the left wings of the Yana specimen (: 220) as '(type ♂)'. I interpret this as a designation of holotype, although the word holotype is not used.

sita Campion (ssp. of *Indothemis limbata*), 1923 : 29-31, fig. Holotype ♂. Ceylon, *Yerbury/Trincomali*, 6.xii.90/*Indothemis limbata sita* Campion, ♂ Holotype. Determined by H. Campion.

smithi Fraser (*Crocorthetrum*), 1921. Kimmins, 1966 : 213.

smithi Kirby (*Cannacria*), 1894b : 266-267. LECTOTYPE ♂. [St. Vincent], Grand Sable (Windward), by pools near sea, Jan. 4/*smithii* [WFK]/*Cannacria smithi* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Brachymesia furcata* Hagen.

stigmata Kirby (*Bradinopyga*), 1893 : 553-554, pl. 41, fig. 3. LECTOTYPE ♂. Ceylon, Trincomalee, 5.viii.90, *Yerbury/stigmaticum* [WFK]/*Bradinopyga stigmata* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

stigmatizans Fabricius (*Libellula*), 1775 : 421. Campion, 1917 : 443.

strachani Kirby (*Apeleutherus*), 1900 : 74-75, pl. 2, figs. 4, 4a. LECTOTYPE ♂. Lagos, Dr. H. Strachan/*Apeleutherus strachani* ♂/App. figd [last two labels in Kirby's writing].

Of the long series recorded by Kirby, only the lectotype ♂ and allotype ♀ bear Kirby's determination labels. I have been unable to trace the ♂ from Free Town, 24.viii, listed by Kirby. This taxon is currently placed as *Bradinopyga strachani* (Kirby).

subpruinusum Kirby (*Sympetrum*), 1886 : 326, pl. 33, fig. 7. LECTOTYPE ♀. N.W. Ind[ia] [WFK]/*subpruinusum* Kirb. type [WFK]/*Sympetrum subpruinusum* Kirby, ♀ Lectotype, D. E. Kimmins det. 1967.

Currently placed as synonym of *Sympetrum commixtum* Selys.

sylvia Kirby (*Nannothemis*), 1889 : 343-344. Holotype ♀. Beara/N. *sylvia* type [WFK]. Currently placed in the genus *Oligoclada*.

terminalis Kirby (*Lathrecista*), 1889 : 336. LECTOTYPE ♂. W. Borneo/*Lathrecista terminalis* ♂ type [WFK]/*Lathrecista terminalis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a synonym of *Lathrecista asiatica* (Fabr.)

thais Kirby (*Perithemis*), 1889 : 324-325. LECTOTYPE ♂. Para/330/*thais* Kb. type [WFK]/*Perithemis thais* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The female, without locality, referred to by Kirby, has not been traced.

thomasi Longfield (*Urothemis*), 1932 : 34-35. Holotype ♂. S. Arabia, Qara Mts., Ain al Rizab, 250 ft., 31.x.1930, *Bertram Thomas/Urothemis thomasi* Longfield, ♂ Holotype.

Miss Longfield's determination label is missing, and a new one has been provided.

- tibialis** Kirby (*Micrathyria*), 1897 : 610, pl. 13, figs. 5-6. LECTOTYPE ♂. Santarem/tibialis [WFK]/*Micrathyria tibialis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- tillyardi** Fraser (*Palaeothemis*), 1923. Kimmins, 1966 : 216.
- torrida** Kirby (*Pseudomacromia*), 1889 : 340-341, pl. 52, fig. 7. LECTOTYPE ♂. Sierra Leone/*Pseudomacromia torrida* type [WFK]/*Pseudomacromia torrida* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- The allotype ♀ (the only one in Kirby's series) is from Sierra Leone and is very teneral, the head now missing. The remaining four syntypes (all males) from Abyssinia, West Africa, Natal and Tenerife, have been traced and are now labelled as paralectotypes. Kirby's statement (: 341) 'Notwithstanding these differences, I am not at present inclined to treat any of these specimens (all of which are males, the only female being described above) as specifically distinct' is considered as definite inclusion of these specimens in the type-series.
- torrida** Kirby (*Tyriobapta*), 1889 : 338-339, pl. 54, figs. 5-6. LECTOTYPE ♂. W. Borneo/*Tyriobapta torrida* type [WFK]/*Tyriobapta torrida* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- trameiformis** Kirby (*Antidythemis*), 1889 : 315, pl. 51, fig. 4. LECTOTYPE ♂. Para/Carolina Lin. Cab./*Antidythemis trameiformis* ♂ type [WFK]/*Antidythemis trameiformis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- The allotype ♀ bears a label *Dythemis?* *trameiformis* Selys, n. sp. ♀, Amazone?
- translucida** Kirby (*Tramea*), 1889 : 315. Holotype ♀. N. India/*translucida* Kb. type [WFK].
- This taxon is currently placed as a synonym of *Trapezostigma limbata* (Desj.).
- triangularis** Kirby (*Rhyothemis*), 1889 : 319. LECTOTYPE ♂. Sar[awak] 78/Wallace/*triangularis* type [WFK]/*Rhyothemis triangularis* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- The example in the BM Type collection, although bearing a type-label, proved to be an imposter, bearing a label Sandakan, 29.iii.1893 (four years after publication of the description). The specimen selected above as type was found in the general collection. It is presumed that the false type was transferred to the type collection for evacuation in the summer of 1940 and has remained undetected until the preparation of this type list.
- tricolor** Kirby (*Orthetrum*), 1893 : 555-556. Holotype ♂. Kandy, 30.vi.92/Ceylon, *Yerbury*/*Orthetrum tricolor* [WFK].
- trifidum** Kirby (*Acisoma?*), 1889 : 341. Holotype ♂ (lacking head and right hind wing). Congo/*Acisoma ascalaphoides?* [in pencil, writing unknown]/*Acisoma trifida*, type [WFK].
- The spelling of the specific name *trifida* has been corrected to *trifidum*.
- tristrigata** Kirby (*Tetrathemis*), 1889 : 343. Holotype ♀. Gil./Gilolo/T. *tristrigata* type [WFK].
- Currently placed as a synonym of *Tetrathemis irregularis leptoptera* Selys.
- turneri** Kirby (*Rhyothemis*), 1894 : 17. LECTOTYPE ♂. Queensland, *Turner*/Turneri Kb. ♂ [WFK]/*Rhyothemis turneri* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- Currently placed as synonym of *Rhyothemis resplendens* Selys.
- tyleri** Kirby (*Trithemis*), 1899 : 364, pl. 15, fig. 1. LECTOTYPE ♂. La Chorrera, Panama/T. Tyler, Named at sight by W. F. K./*Trithemis tyleri* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
- uniformis** Kirby (*Belonia*), 1889 : 333-334. Holotype ♀. Mex[ico]/*uniformis*, type [WFK].
- Currently placed as a synonym of *Libellula croceipennis* Selys.
- variegatum** Kirby (*Acisoma*), 1898 : 239-240. LECTOTYPE ♂. Fort Johnston, Nyassaland, P. Rendall/*variegatum* Kirby paratype [Distant's writing]/*Acisoma variegatum* Kirby, ♂ Lectotype, D. E. Kimmins det. 1961.
- Currently placed as a synonym of *Acisoma ascalaphoides* Rambur.

venusta Kirby (*Micrathyria*), 1897 : 612-613, pl. 13, fig. 1. Holotype ♂. Macapá, 25.ii.96/Macapa, Lower Amazon, 25.ii.96/venusta [WFK].

This taxon is currently placed in the genus *Erythrodiplax*.

vibrans Fabricius (*Libellula*), 1793 : 380. *Campion*, 1917 : 449-450.

vittata Kirby (*Urothemis*), 1893 : 552, pl. 42, fig. 2. LECTOTYPE ♀. Trincomali, 10.ix.91/Ceylon, *Yerbury/vittata* type [WFK]/*Urothemis vittata* Kirby, ♀ Lectotype, D. E. Kimmins det. 1967.

vittatus Kirby (*Hydrobasileus*), 1889 : 314, pl. 51, fig. 10. Holotype ♂. Men[ado]/Wallace/Tramea 4 vittata type [WFK]/*Hydrobasileus vittatus* Kirby ♂, det. D. E. Kimmins, 1967.

The name *Tramea quadrivittata* Hag. MS is quoted by Kirby in synonymy of *Hydrobasileus vittatus* Kirby.

vivax Calvert (*Brechmorhoga*), 1906 : 279, 280, pl. 8, fig. 46. Holotype ♂. Zapote, Guatemala, *Champion/Brechmorhoga vivax* Calvert, TYPE. P. P. Calvert det. 1906. B. C. A. Neur. p. 280. Original of Pl. viii, f. 46.

wilsoni Pinhey (*Brachythemis*), 1952 : 16, fig. 5A. Holotype ♂. Xambio, S. Sudan, xii. 1950, *C. E. Wilson/Brachythemis* ?sp. nov. E. Pinhey det. 1951/*Brachythemis wilsoni* Pinh. ♂.

woodfordi Kirby (*Nesocria*), 1889 : 335. Holotype ♂. [Solomon Islands], Alu/Nesocria woodfordi type [WFK].

Currently placed in the genus *Protorthemis*.

yerburyi Kirby (*Tetrathemis*), 1893 : 556-557, pl. 41, fig. 4. LECTOTYPE ♂. Kandy, 30.vi.92. B. in coitu A/Ceylon, *Yerbury/yerburii* type [WFK]/*Tetrathemis yerburyi* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.

The specific name, spelt *yerburii* by Kirby, has been emended to *yerburyi*.

Family CORDULIIDAE

aculeata Fraser (*Macromia*), 1927. Kimmins, 1966 : 176.

aeneothorax Nunney (*Ceratogomphus*?), 1895 : 349-351, figs. A, B. Holotype ♂ (one sup. app. and inf. app. glued to card). *Ceratogomphus aeneothorax* Nunney mss, on reverse 'Sierra Leone' and TYPE.

I do not recognise the writing (in red ink) of the locality 'Sierra Leone'. Currently placed in the genus *Macromia*.

annaimallaiensis Fraser (*Macromia*), 1931. Kimmins, 1966 : 178.

atrifrons McLachlan (*Syncordulia*), 1883 : xc-xci. Holotype ♀. Queensland/Type [McL. label]/*Syncordulia atrifrons* McL.

Currently placed in the genus *Micromidia*.

atuberculata Fraser (*Macromia*), 1922. Kimmins, 1966 : 180.

aurolineata Tillyard (ssp. of *Metathemis guttata*), 1913 : 575. LECTOTYPE ♂. Dorrigo, N.S.W., *R. J. Tillyard*, 4.xii.1911/*Metathemis guttata aurolineata* Tillyard, TYPE ♂, R.J.T./*Metathemis guttata aurolineata* Till., ♂ Lectotype, D. E. Kimmins det. 1968.

Currently placed as *Eusynthemis guttata aurolineata* (Tillyard).

bellicosa Fraser (*Macromia*), 1924. Kimmins, 1966 : 181.

berthoudi Tillyard (*Hesperocordulia*), 1911 : 376-378, pl. 10, figs. 1, 3, 9. LECTOTYPE ♂. Waroona, W. A[ustralia], *G. F. Berthoud*, 27.xi.1909/*Hesperocordulia Berthoudi* [sic] Till., ♂ TYPE, R.J.T./*Hesperocordulia berthoudi* Till., ♂ Lectotype, D. E. Kimmins det. 1968.

binocellata Fraser (*Macromia*), 1924. Kimmins, 1966 : 183.

bispina Fraser (*Macromia*), 1954. Kimmins, 1966 : 183.

braueri Selys (*Epithea*), 1871 : 284-285. Holotype ♂. N[ew] Zeal[and]/24/*Cordulia Braueri* de Selys/Type [McL. label]/*Epithea braueri* Selys, ♂ Holotype, D. E. Kimmins det. 1967.

Placed by Fraser, 1939, in his genus *Antipodochlora*.

- burliyarensis*** Fraser (ssp. of *Idionyx corona*), 1922. Kimmins, 1966 : 184.
- canis*** McLachlan (*Tetragoneuria*), 1886 : 104-105. Holotype ♂. [Western North America] Washington Territory [K. H. Morrison]/Type [McL. label]/*Tetragoneuria canis* McL.
- circularis*** Tillyard (*Pseudocordulia*), 1908 : 743, pl. 22, fig. 1. Holotype ♂. Kuranda, N.Q., xii.07, F. P. Dodd/Pseudocordulia circularis Till., ♂ TYPE, R.J.T.
- claviculata*** Tillyard (*Synthemis*), 1908b : 749-750, pl. 22, fig. 5. Holotype ♂. Kuranda, N. Q., i.08, E. Allen/Synthemis claviculata Till., ♂ TYPE, R.J.T.
- contumax*** Selys (*Phyllomacromia*), 1879 : 103-104. Holotype ♂ (right fore wing, both hind wings somewhat damaged). West Africa, Akele/Type [McL. label]/*Phyllomacromia contumax*, Selys [McL. hand].
Currently placed in the genus *Macromia*.
- corona*** Fraser (*Idionyx*), 1921. Kimmins, 1966 : 187.
- corycia*** Laidlaw (*Macromia*), 1922 : 220, 225, fig. 5. Holotype ♂. Sarawak, Ulu Baram 3.xi.1920, J. C. Moulton/Macromia corycia sp. n. ♂ Type, det. F. F. Laidlaw.
The original paper label, in Laidlaw's handwriting, gives the date wrongly as 3.xi.30.
- costalis*** Selys (*Cordulia*), 1871 : 273. Holotype ♀. Georgia/Cordulia costalis de Selys/C. costalis De Selys ♀.
Currently placed as a synonym of *Tetragoneuria cynosura* (Say).
- cupricincta*** Fraser (*Macromia*), 1924. Kimmins, 1966 : 187.
- cupricolor*** Fraser (*Hemicordulia*), 1927. Kimmins, 1966 : 187.
- cyanitincta*** Tillyard (*Synthemis*), 1907 : 724-726, pl. 35, figs. 3-4, pl. 36, fig. 1. LECTOTYPE ♂. Margaret R., W. A., i.07, R. J. Tillyard/Synthemis cyanitincta Till., ♂ TYPE, R.J.T./Synthemis cyanitincta Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.
- cydippe*** Laidlaw (*Macromia*), 1922 : 220-222, 223, figs. 2, 3. Holotype ♂. Sarawak, Lio Matu, 3.x.1914, J. C. Moulton/Macromia cydippe Laidlaw, ♂ Type.
- cyrene*** Lieftinck (*Synthemis*), 1953a : 78-81, figs. 5-9. Holotype ♂. N. E. New Guinea, Saiko, 5,500-6,000 ft., Babu River (Upper Waria River), ix-x.1936/F. Shaw-Mayer/Synthemis cyrene sp. n. ♂.
- divergens*** Tillyard (*Cordulephya*), 1917 : 467-469, pl. 23, figs. 9-10, text-fig. 6. LECTOTYPE ♂. Hornsby, N.S.W., 8.v.16, R. J. Tillyard/Cordulephya divergens Till., ♂ TYPE, R.J.T./Cordulephya divergens Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.
- donaldi*** Fraser (*Indomacromia*), 1924. Kimmins, 1966 : 189.
- elliptica*** Tillyard (*Pseudocordulia*), 1913b : 229-230, pl. 15, figs. 1-2. LECTOTYPE ♂. Kuranda, N. Q[ueensland], F. P. Dodd, 3.i.1913/Pseudocordulia elliptica Tillyd., ♂ TYPE, R.J.T./Pseudocordulia elliptica Till., ♂ Lectotype, D. E. Kimmins det. 1968.
- ellisoni*** Fraser (*Macromia*), 1924. Kimmins, 1966 : 190.
- elongata*** Champion (*Metaphya*), 1921 : 64-66, pl. 9, fig. 17. Holotype ♀. Loc. Baie Ngo, no. 204, 10.ii.14. Coll. P. D. Montague, New Caledonia Exped./New Caledonia, P. D. Montague/Metaphya elongata Champion, Holotype, Determined by H. Champion.
- euterpe*** Laidlaw (*Macromia*), 1915 : 26-29, figs. 1-2. LECTOTYPE ♂. Kina Balu, British North Borneo, J. C. Moulton/M. euterpe Type ♂, 12.ix.13, 7 [Laidlaw's writing]/Macromia euterpe Laidlaw, ♂ Lectotype, D. E. Kimmins det. 1967.
The example chosen as Lectotype has for many years been labelled 'Type H.T.', but was not so designated in print by Laidlaw.
- evelynae*** Lieftinck (*Synthemis*), 1953a : 81-84, figs. 10-11. Holotype ♂ (abd. segs. 6-10 and anal appendages wanting). Papua, Mafulu, 4,000 ft., xii.1933, L. E. Cheesman/Synthemis evelynae sp. n. ♂.

- fallax* McLachlan (*Gomphomacromia*), 1881 : 141-142. LECTOTYPE ♂. Intaj, Ecuador/*Gomphomacromia fallax* McL./Type [McL. label]/*Gomphomacromia fallax* McL., ♂ Lectotype, D. E. Kimmins det. 1967.
- fenella* Campion (*Synthesis*), 1921 : 61-62, pl. 9, fig. 16. text-fig. 11. Holotype ♂. Loc. Mt. Mou, No. 725, Date 20.iii.14. Coll. P. D. Montague, New Caledonia Exped./*Synthemis fenella* Campion, Holotype. Determined by H. Campion.
- fidelis* McLachlan (*Hemicordulia*), 1886 : 104. Holotype ♂. [Loyalty Isls.] Uvea/Type [McL. label]/*Hemicordulia fidelis* McLach.
- flavicauda* McLachlan (*Nesocordulia*), 1882 : clxx-clxxii. Holotype ♂. Madagascar/*Nesocordulia flavicauda* McL./Type [McL. label].
- flavicincta* Kirby (*Phyllomacromia*), 1898 : 241-242. LECTOTYPE ♂. Rustenberg, W. L. D[istant]/*flavicincta* (type) Kirby [Distant's writing]/*Macromia picta* Selys H. C[ampion]/*Phyllomacromia flavicincta* Kirby, ♂ Lectotype, D. E. Kimmins det. 1967.
Currently placed in the genus *Macromia*, where it is a junior homonym of *M. flavicincta* Selys and a synonym of *Macromia picta* (Selys).
- flavicincta* Selys (*Macromia*), 1874 : 25-27. LECTOTYPE ♂. North Ind[ia]/*Macr. flavicincta* de Selys/*Macromia flavicincta* Selys, ♂ Lectotype, D. E. Kimmins det. 1967.
The male chosen as Lectotype has for many years been labelled Holotype, but it was not so indicated in print by Selys. The original description gives 'Nord du Bengale' as locality, but it is only the ♀ which is so labelled.
- flavocolorata* Fraser (*Macromia*), 1922. Kimmins, 1966 : 192.
- flavovittata* Fraser (*Macromia*), 1935. Kimmins, 1966 : 192.
- flexicauda* Campion (*Synthemis*), 1921 : 57-61, pl. 8, fig. 14, pl. 9, fig. 15, text-fig. 10. Holotype ♂. Loc. Mt. Nekando, Date, 24.v.14, Coll. P. D. Montague, New Caledonia Exped./New Caledonia, P. D. Montague/*Synthemis flexicauda* Campion ♂. Holotype. Determined by H. Campion.
- fulva* Laidlaw (*Macromidia*), 1915 : 29-30, fig. 3. Holotype ♂. Borneo, J. C. Moulton/*Macromidia fulva*, ♂ Laidlaw, TYPE. Kina Balu, Borneo, 11.ix.1913, J. C. Moulton.
- fulvia* Fraser (ssp. of *Idionyx corona*), 1924. Kimmins, 1966 : 193.
- galeata* Fraser (*Idionyx*), 1924. Kimmins, 1966 : 193.
- genialis* Laidlaw (*Macromidia*), 1923 : 232. Holotype ♂. Malay Peninsula, Pahang, Gunung Tahan, 10.xii.1921/*Macromidia genialis* ♂ Type [Laidlaw's writing].
- gomphomacromioides* Tillyard (*Synthemioptis*), 1917 : 464-467, pl. 23, figs. 7-8, text-figs 4-5. LECTOTYPE ♂. Cradle Mt., Tas., R. J. Tillyard, 18.i.17/*Synthemioptis gomphomacromioides* Till., ♂ TYPE, R.J.T./*Synthemioptis gomphomacromioides* Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.
- gracillima* Fraser (*Hemicordulia*), 1944. Kimmins, 1966 : 194.
- halei* Fraser (*Macromia*), 1928. Kimmins, 1966 : 194.
- ida* Fraser (*Macromia*), 1924. Kimmins, 1966 : 196.
- imbricata* Fraser (*Idionyx*), 1926. Kimmins, 1966 : 196.
- indica* Fraser (*Macromia*), 1924. Kimmins, 1966 : 197.
- insignis* Kirby (*Pseudogomphus*), 1889a : 299. Holotype ♂. Cameroons/*Pseudogomphus insignis* [WFK].
Currently placed as a synonym of *Macromia sophia* Selys. The tenth abdominal segment and anal appendages are detached.
- insularis* Kimmins (*Guadalca*), 1957a : 318-319, figs. 3-4. LECTOTYPE ♂. Solomon Is., Guadalcanal, Tapenanje, 10-23.xii.1953, J. D. Bradley/*Guadalca insularis* Kim., ♂ Type, D. E. Kimmins det. 1955/*Guadalca insularis* Kimmins, ♂ Lectotype, D. E. Kimmins det. 1967.

- intricata* Fraser (*Idionyx*), 1926. Kimmins, 1966 : 198.
- irata* Fraser (*Macromia*), 1924. Kimmins, 1966 : 198.
- junior* Pinhey (ssp. of *Macromia kimminsi*), 1961 : 116, pl. 8, figs. 10, 15, 16. Holotype ♂. *van Someren*, Broderick Falls, Scarp, Kenya, v. 51/*Macromia kimminsi junior* Pinh. HOLOTYPE.
- kimminsi* Fraser (*Macromia*), 1954. Kimmins, 1966 : 199.
- kimminsi* Lieftinck (*Synthemis*), 1953a : 75-78, figs. 1-4. Holotype ♂. N.E. New Guinea, Saiko, 5,500-6,000 ft., Babu River (Upper Waria River), 24.x.1936/*F. Shaw Mayer/Synthemis kimminsi* sp. n.
- laidlawi* Fraser (*Idionyx*), 1936. Kimmins, 1966 : 200.
In my paper (1966), the reference to this species was omitted. It should be 1936a : 701.
- leachi* Selys (*Synthemis*), 1871a : 561. Holotype ♂. N[ew] Holland/Cord. Leachii ♂ Selys/Synth. Leachii de Selys.
The left hind wing is missing and the other wings are somewhat battered. The specific name has been emended to 'leachi'.
- leoni* Fraser (*Phyllomacromia*), 1928. Kimmins, 1966 : 200.
- lieftincki* Fraser (*Macromia*), 1954. Kimmins, 1966 : 201.
- magnifica* Selys (*Macromia*), 1874 : 22-24. LECTOTYPE ♀. N. California, *Walsingham*/*Macromia magnifica* ♀ [Selys' writing]/*Macromia magnifica* Selys, ♀ Lectotype, D. E. Kimmins det. 1967.
The ♂ syntype of this species has not been traced in the McLachlan collection. The ♀ was found (without any type label) and has been designated Lectotype. Selys gives no indication that he retained the ♂, and Martin (1906 : 67) writes 'Types ♂ et ♀ coll. MacLachlan'.
- malabarensis* Fraser (*Epophthalmia*), 1935. Kimmins, 1966 : 203.
- malayana* Laidlaw (ssp. of *Macromia moorei*), 1928 : 133-134. Holotype ♂. Pahang, L. Tamang, 13.vi.1923, *F. N. Chasen*/*Macromia moorei malayana* n. subsp. [Laidlaw's writing]/*Macromia moorei malayana* ♂ Holotype, Laidlaw [DEK].
Currently placed as synonym of *Macromia moorei fumata* Kruger.
- marginata* Selys (*Somatochlora?*), 1883 : 109-110. Holotype ♂ (head lacking). Japan/Type [McL. label]/*Somatochlora? marginata* Selys, ♂, Japon [Selys' writing].
Currently placed in the genus *Epitheca*.
- [*martini* Tillyard (*Synthemis*), 1907 : 726-728, pls. 35, figs. 1-2 and 36, fig. 2.]
There are no examples of this species in BM(NH) collection. The types may perhaps still be in Canberra.
- mascarenica* Fraser (*Nesocordulia*), 1948. Kimmins, 1966 : 203.
- melanosoma* Tillyard (ssp. of *Metathemis guttata*), 1913b : 231-233. LECTOTYPE ♂. Tambourine, Q[ueensland], *R. J. Tillyard*, 24.xii.1912/*Metathemis guttata melanosoma* Tillyd., ♂ TYPE, R.J.T./*Metathemis guttata melanosoma* Till., ♂ Lectotype, D. E. Kimmins det. 1968.
Currently placed as *Eusynthemis guttata melanosoma* (Tillyard).
- metallica* Tillyard (*Lathrocordulia*), 1911 : 379-380, pl. 10, figs. 2, 4 and 10. LECTOTYPE ♂. Waroona, W. A[ustralia], *G. F. Berthoud*, 3.xii.1910/*Lathrocordulia metallica* Till., ♂ TYPE, R.J.T./*Lathrocordulia metallica* Till., ♂ Lectotype, D. E. Kimmins det. 1968.
The lectotype lacks the apex of the left superior anal appendage.
- micans* Laidlaw (*Metaphya*), 1912 : 2-3, pl. 1. Holotype ♂. Sarawak, 1910/*Metaphya micans* Type ♂ [Laidlaw's writing].
The allotype ♀ was described later (*Proc. zool. Soc. Lond.* 1913 : 65-67).
- minima* Fraser (*Idionyx*), 1931. Kimmins, 1966 : 204.

- montaguei** Campion (*Synthemis*), 1921 : 55-57, pl. 8, fig. 13, text-fig. 9. Holotype ♂. Loc. Alt Mou, Date, 10.iii.14 (No. 488), Coll. P. D. Montague, New Caledonia Exped./New Caledonia, P. D. Montague/Synthemis montaguei Campion, ♂, Holotype. Determined by H. Campion.
- montana** Tillyard (*Cordulephya*), 1911a : 392-397, pls. 11-12. Holotype ♂. [Medlow], Blue Mts., N.S.W., 19.i.10, R. J. Tillyard/Cordulephya montana Till., ♂ TYPE, R.J.T.
Holotype ♂ specified by Tillyard (1911 : 397, line 14, 'type-male'). The locality 'Medlow' is given in the text in several places but not on the locality label.
- moorei** Selys (*Macromia*), 1874 : 28. Holotype ♂. Himalaya/Macromia moorei Selys ♂, n. sp.
- mystica** Tillyard (*Austrophya*), 1908b : 739-740, pl. 31, fig. 1. Holotype ♀. Kuranda, N. Q., i.08, F. P. Dodd/Austrophya mystica Till., TYPE ♀, R.J.T.
- nadganiensis** Fraser (*Idionyx*), 1924. Kimmins, 1966 : 205.
- nasalis** Selys (*Epitheca*), 1874 : 21. Holotype ♀. North America/Epith. nasalis de Selys mss., var. d'albicincta B.? [Selys' writing].
This example has at some time lost its locality label and a new one has been provided. Currently placed in the genus *Somatochlora*.
- nigra** Tillyard (*Synthemis*), 1906 : 489-491. LECTOTYPE ♂ (lacking most of left hind wing). Kuranda, N. Q., i.05, R. J. Tillyard/Synthemis nigra Till. ♂ TYPE, R.J.T./Synthemis nigra Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.
Currently placed in the genus *Eusynthemis*.
- nilgiriensis** Fraser (ssp. of *Idionyx corona*), 1922. Kimmins, 1966 : 206.
- nilgiriensis** Fraser (*Phyllomacromia*), 1918. Kimmins, 1966 : 207.
- occidentalis** Tillyard (ssp. of *Synthemis macrostigma*), 1910 : 355. LECTOTYPE ♂. Bridgetown, W. A., i.07, R. J. Tillyard/Synthemis macrostigma Selys, subsp. occidentalis Till., ♂ TYPE, R.J.T./Synthemis macrostigma occidentalis Till., ♂ Lectotype, D. E. Kimmins det. 1967.
- olivei** Tillyard (*Synthemis*), 1908b : 747-749, pl. 22, fig. 4. LECTOTYPE ♂. Cooktown, N. Q., i.08, R. J. Tillyard/Synthemis olivei Till., ♂ TYPE, R.J.T./Synthemis olivei Tillyard, ♂ Lectotype, D. E. Kimmins det. 1967.
Currently placed in the genus *Choristhemis*.
- orientalis** Tillyard (ssp. of *Synthemis macrostigma*), 1910 : 355. LECTOTYPE ♂. Blue Mts., N.S.W., ii.05, R. J. Tillyard/Synthemis macrostigma Selys, subsp. orientalis Till., ♂ TYPE/Synthemis macrostigma orientalis Till., ♂ Lectotype, D. E. Kimmins det. 1967.
- ornata** Fraser (*Idionyx*), 1921. Kimmins, 1966 : 208.
- pacifica** Fraser (*Hemicordulia*), 1925. Kimmins, 1966 : 208.
- pallida** Fraser (*Macromia*), 1924. Kimmins, 1966 : 208.
- periyashola** Fraser (*Idionyx*), 1939. Kimmins, 1966 : 209.
- piscator** Barnard (*Presba*), 1934 : 168, figs. 1f-h, 2. LECTOTYPE ♂. Groot Drakenstein, Cape Province, xii.1932, A. C. Harrison/Presba piscator Barnard ♂ TYPE [DEK]/Presba piscator Barnard, ♂ Lectotype, D. E. Kimmins det. 1967.
- pseudaficana** Pinhey (*Macromia*), 1961 : 109, pl. 8, fig. 5. Holotype ♂. Aswe R., Karimoja Suk, Uganda, iv.1952/Macromia africana Selys, E. Pinhey det. 1952/Macromia africana Selys, F. C. Fraser det. 1953/Macromia pseudaficana Pinh. HOLOTYPE.
Pinhey states that Fraser's determination was made 'with reservation'.
- pygmaea** Selys (*Cordulephya*), 1871 : 316. Holotype ♂. Melbourne, Edwards/122/Cordulephya pygmaea Selys ♂/Type [McL. label].

- refracta* Tillyard (*Austrocordulia*), 1908b : 744-745, pl. 22, fig. 2. Holotype ♂. Cooktown, N. Q., i. 08, *R. J. Tillyard/Austrocordulia refracta* Till., ♂ TYPE, R.J.T.
- regina* Selys (*Synthemis*), 1874 : 31-33. LECTOTYPE ♂. N[ew] S[outh] W[ales]/Synth. regina de Selys/Synthemis regina Selys, ♂ Lectotype, D. E. Kimmins det. 1967.
Selys gives as locality 'Queensland', but the BM(NH) specimen, which he lists first, is labelled New South Wales, the MacLachlan examples being from Queensland. Currently placed in the genus *Eusynthemis*.
- rhinoceroides* Fraser (*Idionyx*), 1934. Kimmins, 1966 : 211.
- rutherfordi* Selys (*Neophya*), 1881 : xvii-xviii, 1 fig. Holotype ♂. Old Calabar, R[utherford]/Type[McL. label]/*Neophya Rutherfordi* Selys.
- saffronata* Fraser (*Idionyx*), 1924, Kimmins, 1966 : 212.
- selysi* Fraser (*Idionyx*), 1926 : 201-202, pls. 9, fig. 6 and 10, fig. 6. LECTOTYPE ♂. I. selysi ♂. Maymyo, Upper Burma, 15.vi.24. Col. *F. Wall* coll. [F.C.F.]/*Idionyx selysi* Fraser. ♂ Lectotype, D. E. Kimmins det. 1967.
Through an unfortunate oversight, this species was omitted from my 1966 paper. The above example has been chosen as Lectotype, in preference to the example in the Selys collection, which Fraser appears not to have seen.
- selysi* Kirby (*Macromia*), 1900 : 77-78. Holotype ♂. Sierra Leone, Free Town, 16.ix.99, *E. E. Austen*/M. Selysi type [WFK].
- shanensis* Fraser (*Macromidia*), 1927. Kimmins, 1966 : 213.
- smithi* White (*Cordulia*), 1845, pl. 6, fig. 2; Selys, 1871 : 261. Holotype ♀. New Zealand/Cordulia Smithii White, Zool. Ereb. & Terror, t. . . , f. . . [A. White's writing].
The first reference to this species is the plate reference given above. The first description is that of Selys, 1871. Currently placed in the genus *Procordulia*.
- sophia* Selys (*Macromia*), 1871a : 550-551. Holotype ♂. Cape Coast Castle [Guinea]/124/Type [McL. label]/*Macromia sophia* ♂ Selys [Selys' writing].
- spiniger* Tillyard (*Synthemis*), 1913 : 572-574, pl. 62, figs. 15-16. Holotype ♂. Waroona, W. A[ustralia], *G. F. Berthoud*, 27.i.1912/*Synthemis spinifer* [sic] Tillyard, TYPE ♂, R.J.T./*Synthemis spiniger* Till., ♂ Holotype, D. E. Kimmins det. 1968.
The specific name was published as 'spiniger' and used consistently five times in the original publication.
- stevensi* Fraser (*Idionyx*), 1924. Kimmins, 1966 : 214.
- subjuncta* Tillyard (ssp. of *Metathemis brevistyla*), 1913 : 574, pl. 62, fig. 1b. LECTOTYPE ♂. Dorrigo, N.S.W., *R. J. Tillyard*, 6.xii.1911/*Metathemis brevistyla subjuncta* Tillyard, TYPE ♂, R.J.T./*Metathemis brevistyla subjuncta* Till., ♂ Lectotype, D. E. Kimmins det. 1968.
Currently placed as *Eusynthemis brevistyla subjuncta* (Tillyard).
- superba* Tillyard (*Hemicordulia*), 1911 : 368-370, pl. 10, figs. 5-8. LECTOTYPE ♂. Pallal, N.S.W., *R. J. Tillyard*, 13.xii.1910/*Hemicordulia superba* Till., ♂ TYPE, R.J.T./*Hemicordulia superba* Till., ♂ Lectotype, D. E. Kimmins det. 1968.
- sylvatica* Fraser (*Macromia*), 1954. Kimmins, 1966 : 215.
- tasmanica* Tillyard (*Synthemis*), 1910 : 344-347, pls. 4, fig. 2 and 7, figs. 2, 20. LECTOTYPE ♂. St. Patrick's R., Tas., 7.i.09, *R. J. Tillyard/Synthemis tasmanica* Till., ♂ TYPE, R.J.T./*Synthemis tasmanica* Till., ♂ Lectotype, D. E. Kimmins det. 1967.
- thalia* Lieftinck (*Macromia*), 1929 : 103-104, fig. 29. Holotype ♂. Khasia Hills/*Macromia thalia* nov. nom., ♂ Holotype, vid. Lieftinck, '28 [Lieftinck's writing]/*Macromia thalia* Lieftinck, ♂[DEK].

This specimen was originally determined as *Macromia ?fraenata* Laidlaw, by Laidlaw.

thetis Ris (*Macromia*), 1921 : 379-381, figs. 65-66. LECTOTYPE ♂. Chirinda For., Gazal[and], 3,600 ft., 9.x.1905, *G. A. K. Marshall*/Macromia spec. a, Det. Dr. F. Ris, Phylomacromia aequatorialis Martin?/Macromia thetis Ris ♂, det. Miss C. Longfield, 1934./Macromia thetis Ris, ♂ Lectotype, D. E. Kimmins det. 1968.

With this type and the allotype ♀ is a mss. note by Miss Longfield. 'These are believed to be the types of *Macromia thetis* Ris, described 1921, in Ann. S. African Mus. Found with all labels as at present, except type labels and determination labels of Miss C. Longfield'.

The female in the South African Museum was excluded from the type series by Ris' reference (p. 381) to the 'typical pair in the British Museum'.

tillyardi Martin (*Macromia*), 1906 : 72. Holotype ♀. Kuranda, i.05/Macromia tillyardi Martin, ♀ TYPE, R.J.T.

Although Tillyard (1906 : 491) records the capture of three females, only one, 'the type-specimen' was sent to Martin and is therefore automatically the holotype.

travencorensis Fraser (*Idionyx*), 1931. Kimmins, 1966 : 216.

trituberculata Fraser (*Macromia*), 1921. Kimmins, 1966 : 216.

unguiculata Fraser (*Idionyx*), 1926. Kimmins, 1966 : 217.

venator Barnard (*Presba*), 1933 : 167, figs. 1a-e. LECTOTYPE ♂. French Hoek, C[ape] P[rovince], with on reverse, K. H. Barnard, xii.1932/Presba venator Barnard, ♂ TYPE [DEK]/Presba venator Barnard, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed as a synonym of *Syncordulia gracilis* (Burm.).

virgula Selys (*Synthemis*), 1874 : 33-34. LECTOTYPE ♂. S. Austral[ia], Bakewell/Synth. virgula de Selys/Synthemis virgula Selys, ♂ Lectotype, D. E. Kimmins det. 1967.

Currently placed in the genus *Eusynthemis*.

viridescens Tillyard (*Macromia*), 1911 : 380-381, pl. 10, fig. 11. Holotype ♂. Cape York, N. Q[ueensland], *H. Elgner*, 22.xi.1909/Macromia viridescens Till., ♀ TYPE, R.J.T./This is said to be the ♀ of *M. terpsichore* Foerster. Tillyd., Proc. Linn. Soc. N.S.W., Vol. xxxvii, 1912, p. 584' [F. C. Fraser's writing].

Currently placed as a synonym of *Macromia terpsichore* Foerster.

whitei Selys (*Macromia*), 1871a : 555. LECTOTYPE ♂. N. India/Macr. Whitei ♂ De Selys [Selys' writing]/Macromia whitei Selys, ♂ Lectotype, D. E. Kimmins det. 1967.

The BM(NH) collection contains 1 ♂, 1 ♀ of this species, labelled by Selys in writing very similar to that on the label of *Synthemis leachi*, described in the same publication. Selys quotes an unspecified number of examples in 'Coll. Selys et Brit. Mus.' Our examples are labelled N. India, which would certainly include part of Bengal. The species is almost certainly named after Dr. Adam White, a previous Keeper of the Zoological Department of the British Museum and one assumes therefore that our examples would be the types. Martin (1906 : 70-71) mentions types of *M. cingulata* Rambur (of which *whitei* is a synonym) in the Selys Collection, but not of *whitei*. I do not know of any previous designation of a type for this species and have therefore selected our ♂ (as above) as Lectotype.

wollastoni Champion (*Synthemis*), 1915 : 486-488, figs. 38-39. Holotype ♀. Dutch New Guinea, *A. F. R. Wollaston*/Snow Mts. 4,000 to 6,000 ft., i-ii.1913/Synthemis wollastoni Cmpn., ♀ Holotype, Determined by H. Champion.

The description gives the additional data, Utakwa River, not mentioned on the printed labels.

xanthosticta Tillyard (ssp. of *Metathemis nigra*), 1913b : 230-231. LECTOTYPE ♂. Tambourine, Q[ueensland], *R. J. Tillyard*, 5.i.1913/Metathemis nigra xanthosticta Till., ♂ Lectotype, D. E. Kimmins det. 1968.

Currently placed as *Eusynthemis nigra xanthosticta* (Tillyard).

zeylanica Fraser (*Macromia*), 1927. Kimmins, 1966 : 219.

REFERENCES

- BARNARD, K. H. 1933. A new genus of Corduline Dragonfly from South Africa (Odonata). *Stylops* **2** (7) : 165-168, figs. 1-2.
- CALVERT, P. P. 1901-1908. Odonata in *Biologia Centrali-Americana*: 17-420, pls. 2-10. London.
- CAMPION, H. 1915. Report on the Odonata collected by the British Ornithologists' Union Expedition and the Wollaston Expedition in Dutch New Guinea. *Trans. zool. Soc. Lond.* **20** (15) : 485-492, text-figs. 38-40.
- 1917. On Fabricius's Types of Odonata in the British Museum (Natural History). *Ann. Mag. nat. Hist.* (8) **19** : 441-450.
- 1921. Odonata collected in New Caledonia by the late Mr. Paul D. Montague. *Ann. Mag. nat. Hist.* (9) **8** : 33-67, pls. 8-9, text-figs. 1-11.
- 1923. A new form of *Indothemis limbaia* from Ceylon (Odonata). *Ann. Mag. nat. Hist.* (9) **11** : 28-31, 1 text-fig.
- FABRICIUS, J. C. 1775. *Systema Entomologiae sistens Insectorum Classes, Ordines, Genera, Species*, etc. pp. 420-426. Flensburgi et Lipsiae.
- 1781. *Species Insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphos*, etc., **1** : 519-528. Hamburgi et Kilonii.
- 1787. *Mantissa Insectorum* etc. **1**. Hafniae.
- 1793. *Entomologia Systematica emendata et aucta*, **2** : viii + 519. Hafniae.
- FRASER, F. C. 1926. A revision of the genus *Idionyx* Selys. *Rec. Indian Mus.* **28** : 195-207, pls. 8-10.
- GARDNER, A. E. 1958. Aspects of the fauna and flora of the Azores. xii(a). Odonata. *Ann. Mag. nat. Hist.* (13) **1** : 791-792.
- KIMMINS, D. E. 1936. Odonata, Ephemeroptera and Neuroptera of the New Hebrides and Banks Island. *Ann. Mag. nat. Hist.* (10) **18** : 68-88, 11 text-figs., pl. 3.
- 1936a. The Odonata of the Oxford University Sarawak Expedition. *J. fed. Malay St. Mus.* **18** (1) : 65-108, 17 text-figs.
- 1942. The type of *Allorhizucha campioni* Ris (Odonata, Libellulinae). *Entomologist* **75** : 47.
- 1943. A new South American Dragonfly. *Ann. Mag. nat. Hist.* (11) **10** : 156-159, figs. 1-6.
- 1953. Miss L. E. Cheesman's Expedition to New Caledonia, 1949—Orders Odonata, Ephemeroptera, Neuroptera and Trichoptera. *Ann. Mag. nat. Hist.* (12) **6** : 241-257, 24 text-figs.
- 1957. Some notes on nomenclature in the family Libellulidae (Odonata). *Entomologist* **90** : 95-97.
- 1958. New species and subspecies of Odonata. *Bull. Br. Mus. nat. Hist. (Ent.)* **7** (7) : 347-358, 7 text-figs.
- 1966. A list of the Odonata Types described by F. C. Fraser, now in the British Museum (Natural History). *Bull. Br. Mus. nat. Hist. (Ent.)* **18** (6) : 173-227.
- KIRBY, W. F. 1884. On the Neuroptera collected during the recent expedition of H.M.S. *Challenger*. *Ann. Mag. nat. Hist.* (5) **13** : 453-456.
- 1886. On a small collection of Dragonflies from Murree and Campbellpore (N.W. India), received from Major J. W. Yerbury, R.A. *Proc. zool. Soc. Lond.* **1886** : 325-329, pl. 33.
- 1889. A revision of the subfamily Libellulinae, with descriptions of new genera and species. *Trans. zool. Soc. Lond.* **12** (9) : 249-348, pls. 51-57.
- 1889a. Descriptions of new genera and species of Odonata in the collection of the British Museum, chiefly from Africa. *Proc. zool. Soc. Lond.* **1889** : 297-303.
- 1890. Description of a new species of Dragonfly. *Ann. Mag. nat. Hist.* (6) **5** : 112-113.
- 1893. Catalogue of the described Neuroptera Odonata (Dragonflies) of Ceylon, with descriptions of new species. *J. Linn. Soc., Zool.* **24** : 545-566, pls. 41-42.
- 1894. On a small collection of Odonata (Dragonflies) from Queensland, with descriptions of five new species. *Ann. Mag. nat. Hist.* (6) **14** : 15-21.

- KIRBY, W. F. 1894a. Descriptions of a new species of Dragonfly (*Dythemis Broadwayi*) from Trinidad. *Ann. Mag. nat. Hist.* (6) **14** : 227-228.
- 1894b. On some small collections of Odonata (Dragonflies) recently received from the West Indies. *Ann. Mag. nat. Hist.* (6) **14** : 261-269.
- 1894c. Notes on a small collection of Odonata etc. from Upper Burma, with the description of a new species. *Ann. Mag. nat. Hist.* (6) **14** : 111-113.
- 1896. On some Dragonflies obtained by Mr. and Mrs. Lort-Phillips in Somaliland. *Proc. zool. Soc. Lond.* **1898** : 521-523.
- 1897. List of the Neuroptera collected by Mr. E. E. Austen on the Amazons etc. during the recent expedition of Messrs. Siemens Bros. Cable S.S. *Faraday*, with descriptions of several new species of Odonata (Dragonflies). *Ann. Mag. nat. Hist.* (6) **19** : 598-617, pls. 12-13.
- 1898. On a collection of Dragonflies from the Transvaal and Nyasaland. *Ann. Mag. nat. Hist.* (7) **2** : 229-245.
- 1899. On a collection of Odonata (Dragonflies) from Panama. *Ann. Mag. nat. Hist.* (7) **3** : 362-371, pl. 15.
- 1900. Report on the Neuroptera Odonata collected by Mr. E. E. Austen at Sierra Leone during August and September 1899. *Ann. Mag. nat. Hist.* (7) **6** : 67-79, pl. 2 and 2 text figs.
- 1900a. On a small collection of Odonata (Dragonflies) from Hainan, collected by the late John Whitehead. *Ann. Mag. nat. Hist.* (7) **5** : 530-539, pl. 12.
- 1905. List of a small collection of Odonata (Dragonflies) from Ceylon, collected by Mr. E. Ernest Green, with notes on the genus *Zygonidia* and its allies, and descriptions of new species of *Zygonidia* Kirb, and *Onychothemis* Brauer, from Ceylon and Tonkin. *Ann. Mag. nat. Hist.* (7) **15** : 270-278.
- 1905a. List of a collection of Neuroptera Odonata (Dragonflies) formed by G. A. K. Marshall, Esq., at Salisbury, Mashonaland, with descriptions of a new genus and two new species. *Ann. Mag. nat. Hist.* (7) **15** : 190-193.
- 1909. Ruwenzori Expedition Reports. 7. Neuroptera. *Trans. zool. Soc. Lond.* **19** (1) : 59-62.
- LAIDLAW, F. F. 1912. On a new genus and species of Odonata from Sarawak. *Sarawak Mus. J.* **1** (2) : 1-3, pl. 1.
- 1915. Contributions to a study of the Dragonfly fauna of Borneo.—Part III. A collection made on Mount Kina Balu by Mr. J. C. Moulton in September and October, 1913. *Proc. zool. Soc. Lond.* **1915** : 23-39, text-figs. 1-5.
- 1922. Notes on some oriental Dragonflies: the genus *Macromia*. *Journ. Straits Br. R. Asiat. Soc.* **85** : 218-229, 7 figs.
- 1923. A survey of the dragonfly fauna of the Malay Peninsula, with notes on that of neighbouring countries. Part I. *J. Malayan Br. R. Asiat. Soc.* **1** : 319-333, figs.
- 1928. Notes on Oriental Dragonflies (Odonata), with descriptions of new species. *Proc. zool. Soc. Lond.* **1928** : 129-138, 3 text-figs.
- LIEFTINCK, M. A. 1929. A revision of the known malaysian Dragonflies of the genus *Macromia* Rambur, with comparative notes on species from neighbouring countries and descriptions of new species. *Tijdschr. Ent.* **72** : 59-108, 22 figs.
- 1953. Revisional notes on the genera *Diplacina* Brauer and *Huonia* Förster (Odon.), *Treubia* **22** (1) : 153-216, 23 text-figs.
- 1953a. Revision of the Australasian species of *Synthemis* Selys (Odon. Corduliidae), with descriptions of four new species and a key to their identification. *Idea* **9** (3-4) : 70-88, 14 figs.
- 1955. Two new species of *Lanthanusa* Ris, from the high mountains of New Guinea (Odonata). *Zoöl. Meded. Leiden* **33** (18) : 157-164, 9 text-figs.
- LONGFIELD, C. 1932. A new species of the genus *Urothemis* from Southern Arabia, and some comments on the species of Odonata inhabiting the Qara Mountains. *Stylops* **1** : 34-35.
- 1936. Studies on African Odonata, with synonymy and descriptions of new species and subspecies. *Trans. R. ent. Soc. Lond.* **85** : 467-498, 10 text-figs.

- LONGFIELD, C. 1947. The Odonata of South Angola. *Archos Mus. Bocage* **16** : 1-31, 11 figs.
 — 1953. The *Atoconeura* problem (Odon., Libellulidae). *Entomologist* **86** : 42-49, pl. 1.
 — 1955. The Odonata of North Angola. Part I. *Publ ções cult. Co. Diam. Angola* **27** (11) : 13-63, 10 figs.
 — 1959. The Odonata of North Angola. Part II. *Publ ções cult. Co. Diam. Angola* **45** : 13-42, 6 text-figs.
- MCLACHLAN, R. 1881. Description of a new species of Corduliina (*Gomphomacromia fallax*) from Ecuador. *Trans. ent. Soc. Lond.* **1881** : 141-142.
 — 1883. Neuroptera of the Hawaiian Islands. Part I. Pseudo-Neuroptera. *Ann. Mag. nat. Hist.* (5) **12** : 226-240.
 — 1883a. Description d'une espèce nouvelle de Corduline du sousgenre *Syncordulia*. *C.r. Soc. ent. Belg.* **27** : xc-xci.
 — 1886. Two new species of Cordulina. *Entomologists' mon. Mag.* **23** : 104-105.
 — 1892. Supplementary note on the Neuroptera of the Hawaiian Islands. *Ann. Mag. nat. Hist.* (6) **10** : 176-179.
 — 1894. On two small collections of Neuroptera from Ta-chien-lu, in the province of Szechuen, Western China, on the frontier of Thibet. *Ann. Mag. nat. Hist.* (6) **13** : 412-436.
 — 1894a. Some additions to the neuropterous fauna of New Zealand, with notes on certain described species. *Entomologists' mon. Mag.* (2) **5** : 238-243, 270-272.
- NUNNEY, W. H. 1895. A new West-African Insect. *Ann. Mag. nat. Hist.* (6) **16** : 349-351.
- PINHEY, E. C. G. 1952. Three new species of Odonata from Eastern Africa. *Occ. Pap. Coryndon meml Mus.* **3** : 13-16, fig. 5.
 — 1955. Some dragonflies of East and Central Africa and a rarity from Mauritius. *Occ. Pap. Coryndon meml Mus.* **4** : 17-41, pls 2-3, 9 text-figs.
 — 1958. Records of Dragonflies from the Zambezi and Rhodesia; a revision of the genus *Platycypha*; a gynandromorph dragonfly from Uganda. *Occ. Pap. natn. Mus. Sth Rhodesia* **22B** : 97-116, 7 text-figs.
 — 1961. *A Survey of the Dragonflies (Order Odonata) of Eastern Africa*. viii + 214 pp., 11 pls., 2 text-figs. British Museum (Nat. Hist.), London.
- RIS, F. 1909-1919. Libellulinae, Catalogue systématique et descriptif. *Coll. zool. Selys-Longchamps*, Bruxelles, fasc. **IX-XVI**, 1, 278 pp., 8 pls., 692 text-figs.
 — 1915a. New Dragonflies (Odonata) of the subfamily Libellulinae from Sierra Leone, W. Africa. *Ann. Mag. nat. Hist.* (8) **15** : 213-223.
 — 1921. The Odonata or Dragonflies of South Africa. *Ann. S. Afr. Mus.* **19** : 245-452, pls. 5-12, 77 text-figs.
- SELYS-LONGCHAMPS, E. DE. 1871. Synopsis des Cordulines. *Bull. Acad. r. Sci. Belg.* (2) **31** : 238-316.
 — 1871a. Synopsis des Cordulines (Suite). Deuxieme Légion.—*Macromia*. *Bull. Acad. r. Sci. Belg.* (2) **31** : 519-565.
 — 1872. Note sur plusieurs Odonates de Madagascar et des Îles mascareignes. *Rev. Mag. Zool.* (2) **23** : 175-183.
 — 1874. Additions au Synopsis des Cordulines. *Bull. Acad. r. Sci. Belg.* (2) **37** : 16-34.
 — 1878. Considérations sur la faune de la Nouvelle Guinée, des Moluques et de la Célèbes. *Mitt. K. zool. mus. Dresden* **1878** : 289-322.
 — 1879. Description of a new species of *Phyllomacromia* (Corduliina) from West Africa. *Entomologists' mon. Mag.* **16** : 103-104.
 — 1881. *Neophya* Selys, nouveau genre de Cordulines. *C.r. Soc. ent. Belg.* **25** : xv-xviii, 1 fig.
 — 1883. Les Odonates du Japon. *Ann. Soc. ent. Belg.* **27** : 82-143.
 — 1887. Revision des Odonates de l'Asie septentrionale, du Japon et de l'Afrique septentrionale. *Ann. Soc. ent. Belg.* **31** : 50-68.
 — 1888. Odonates recueillis aux Îles Loo-Choo par feu M. Pryer. *C.r. Soc. ent. Belg.* (3) **101** : xlviiii-liiii.
- TILLYARD, R. J. 1906. New Australian species of the family Libellulidae. *Proc. Linn. Soc. N.S.W.* **31** : 480-492, pl. 44.

- TILLYARD, R. J. 1907. The Dragonflies of South-western Australia. *Proc. Linn. Soc. N.S.W.* **32** : 719-742, pls. 34-36.
- 1908. On some remarkable Australian Libellulinae. *Proc. Linn. Soc. N.S.W.* **33** : 637-649, pl. 14.
- 1908a. On the genus *Nannodythemis*, with descriptions of new species. *Proc. Linn. Soc. N.S.W.* **33** : 444-445, pl. 6.
- 1908b. On some remarkable Australian Corduliinae, with descriptions of new species. *Proc. Linn. Soc. N.S.W.* **33** : 737-751, pls. 21-22.
- 1910. Monograph of the genus *Synthemis*. *Proc. Linn. Soc. N.S.W.* **35** : 312-377, pls. 4-9.
- 1911. Further notes on some rare Australian Corduliinae, with descriptions of new species. *Proc. Linn. Soc. N.S.W.* **36** : 366-387, pl. 10.
- 1911a. On the genus *Cordulephya*. *Proc. Linn. Soc. N.S.W.* **36** : 388-422, pls. 11-12.
- 1913. On some Australian Anisoptera, with descriptions of new species. *Proc. Linn. Soc. N.S.W.* **37** : 572-584, pl. 64.
- 1913a. Description and life history of a new species of *Nannophlebia*. *Proc. Linn. Soc. N.S.W.* **37** : 712-726, pl. 74.
- 1913b. Some descriptions of new forms of Australian Odonata. *Proc. Linn. Soc. N.S.W.* **38** : 229-241, pl. 15.
- 1917. On some new dragonflies from Australia and Tasmania [Order Odonata]. *Proc. Linn. Soc. N.S.W.* **42** : 450-479, pl. 23 and 10 text-figs.
- WHITE, A. 1846-1874. In *The Zoology of the Voyage of H.M.S. Erebus and Terror, under the command of Captain Sir James Clark Ross, R.N., F.R.S., during the years 1839 to 1843*. Ed. Richardson, J. & Gray, J. E. **2**. Insects, p. 25. London.

